

Raptor[®] herbicide

Technical Information Bulletin for the Use of Raptor[®] Herbicide to Aid in the Control of Volunteer Canola in Roundup Ready[®] Soybeans in Minnesota

EPA Reg. No. 241-379

This recommendation is made as permitted under FIFRA section 2 (ee) and had not been submitted to or approved by the EPA.

All applicable directions, restrictions, precautions, and Conditions of Sale and Warranty on the EPA label are to be followed. Refer to the container label for additional instructions. Always read and follow label directions.

Information contained in this Technical Information Bulletin is not intended to replace or amend any product labeling. Always read and follow all label directions when using any pesticide alone or in mix combinations.

The user must have this recommendation in their possession at the time of use.

This recommendation is for the use of Raptor herbicide to aid in control of volunteer canola in soybeans in Minnesota.

DIRECTIONS FOR USE:

Apply **Raptor** to emerged volunteer canola in soybeans but prior to soybean bloom stage. Use 0.016 lb imazamox a.e./acre (2 fluid ounce of **Raptor** per acre) when tankmixed with glyphosate in Roundup Ready soybeans. Use the appropriate surfactants as recommended by the glyphosate label. **Raptor** will aid in the control of volunteer canola (Roundup Ready and conventional) from 1-8 inches tall.

Raptor will not control volunteer **Clearfield**^{*} Canola.

Raptor and **Clearfield** are registered trademarks of BASF.
Roundup Ready is a registered trademark of Monsanto.

© 2004 BASF Corporation
All rights reserved

00241-00379.2004-0309.NVA **2004-04-133-0049**

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709