

# Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## CleanWave<sup>®</sup> Herbicide

EPA Reg. No. 62719-525

**For Control Of Annual And Perennial Broadleaf Weeds In Wheat (Including Durum)**

**For Use Only in Montana**

### ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for CleanWave<sup>®</sup> herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of CleanWave according to this supplemental labeling is subject to all use precautions, restrictions and limitations imposed by the label affixed to the container.

### Directions for Use

#### Tank Mixtures for Wheat (including Durum)

CleanWave<sup>®</sup> herbicide may be applied in tank mix combination with labeled rates of other products registered for postemergence application in wheat. See Tank Mixing Precautions under Mixing Instructions. When tank mixing, do not exceed specified application rates and use only in accordance with the most restrictive precautions and limitations on the respective product labels. When tank mixing with 2,4-D ester, users should make certain that the crop has reached the minimum growth stage for application (4-leaf with 3-4 tillers). Applications of 2,4-D ester made prior to the 4-leaf, tillering stage of growth can result in crop injury.

The following broadleaf and grass herbicide tank mix partners are specified to improve control of certain weed species:

Tank Mix Partner	Rate/Acre	Weed Species
2,4-D ester	1/2 to 3/4 pint <sup>†</sup>	mustard, pigweed, Canada thistle, Russian thistle, field bindweed
MCPA ester or amine	1/2 to 3/4 pint <sup>†</sup>	mustard
Harmony <sup>®</sup> GT herbicide	1/12 to 3/10 oz	mustard, pigweed, Russian thistle
Express <sup>®</sup> XP herbicide	1/8 to 1/3 oz	mustard, Canada thistle, Russian thistle
Ally <sup>®</sup> XP herbicide	1/10 oz	mustard, pigweed, Russian thistle
Axial <sup>™</sup> Axial XL	8.2 fl oz 16.4 fl oz	wild oats, green foxtail, yellow foxtail, Persian dandel
Axial+MCPA ester	8.2 fl oz+1/2 to 3/4 pint <sup>†</sup>	wild oats, green foxtail, yellow foxtail, Persian

Axial XL+MCPA ester	16.4 fl oz+1/2 to 3/4 pint <sup>†</sup>	darnel, other broadleaf weeds
Discover <sup>®</sup>	3.2-4 fl oz	wild oats, green foxtail, yellow foxtail, Persian darnel
Discover NG	12.8-16 fl oz	
Discover+MCPA Discover NG+MCPA	3.2-4 fl oz or 3.2-4 fl oz+1/2 to 3/4 pint <sup>†</sup> 12.8-16 fl oz or 12.8-16 fl oz+1/2 to 3/4 pint <sup>†</sup>	wild oats, green foxtail, yellow foxtail, Persian darnel, other broadleaf weeds
Olympus <sup>™</sup>	0.6 - 0.9 oz	Brome species, cheat
Olympus+MCPA amine	0.6 - 0.9 oz+1/2 to 3/4 pint <sup>†</sup>	Brome species, cheat, other broadleaf weeds
Olympus Flex	3-3.5 oz	Brome species, cheat, Italian ryegrass
Olympus Flex+MCPA amine	3-3.5 oz+1/2 to 3/4 pint <sup>†</sup>	Brome species, cheat, Italian ryegrass, other broadleaf weeds
PowerFlex <sup>™</sup>	3.5 oz	Brome species, cheat, Italian ryegrass, wild oat
PowerFlex+MCPA	3.5 oz+1/2 to 3/4 pint <sup>†</sup>	Brome species, cheat, Italian ryegrass, wild oat, other broadleaf weeds

<sup>†</sup>Rate based on a 4 lb/gal formulation

**Restrictions:**

- Do not tank mix CleanWave with Osprey<sup>™</sup> or Osprey+MCPA
- There are no restrictions on grazing following application of CleanWave at labeled rates.
- Do not apply more than 14 fl oz per acre of CleanWave per growing season.
- Do not apply more than once per growing season.
- **Preharvest Interval:** Do not apply within 14 days of harvest for hay or within 50 days of harvest for grain and straw.

<sup>®</sup>Trademark of Dow AgroSciences LLC

R328-012  
EPA accepted 04/27/10  
Replaces: Initial