



SUPPLEMENTAL LABELING

AQUA STAR

42750-59

READ THE ENTIRE LABEL FOR AQUA STAR BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL. THIS SUPPLEMENTAL LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SILVICULTURAL SITES AND UTILITY RIGHTS-OF-WAY

TYPES OF APPLICATIONS: This product is recommended for the control or partial control of woody brush, trees and herbaceous weeds. This product is labeled for use in forestry and utility sites. This product is also recommended for use in preparing or establishing wildlife openings within these sites and maintaining logging roads, and for side trimming along utility rights-of-way.

In forestry, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

In utilities, this product is recommended for use along electrical power, pipeline and telephone rights-of-way, and in other utility sites associated with these rights-of-way, such as substations.

APPLICATION RATES AND TIMING:

Application	AQUA STAR	Spray Volume (Gal/A)
Broadcast		
Aerial	1.5 to 7.5 qts/a	5 to 30
Ground	1.5 to 7.5 qts/a	10 to 60
Spray-to-Wet		
Handgun, Backpack, Mistblower	0.6% to 2% by volume	spray-to-wet
Low Volume Directed Spray		
Handgun, Backpack, Mistblower	4% to 7.5% by volume	partial coverage*

*For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the plant is important for best results.

In forestry site preparation and utility rights-of-way applications, this product requires use with a nonionic surfactant. Use a nonionic surfactant with greater than 80 percent active ingredient and labeled for use with herbicides. Use of this product without surfactant will result in reduced performance. See the "MIXING" section of this labeling for more information.

Mix 2 or more quarts of the nonionic surfactant per 100 gallons of spray solution (0.5 percent or more by spray volume). Use of surfactant concentrations greater than 1.5 percent by spray volume with handgun applications or 2.5 percent by spray volume with broadcast applications is not recommended.

Use higher rates of this product within the recommended range for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Increase rates within the recommended range for control of perennial herbaceous weeds any time after emergence and before seedheads, flowers or berries appear.

Use the lower rates of this product within the recommended range for control of annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to the foliage of actively growing annual herbaceous weeds any time after emergence.

This product has no herbicidal or residual activity in the soil. Where repeat applications are necessary, do not exceed 8 quarts of this product per acre per year.

Tank Mixtures

Tank mixtures of this product may be used to increase the spectrum of vegetation controlled. When tank mixing, read and carefully observe the label claims, cautionary statements and all information on the labels of both products used. Use according to the most restrictive precautionary statements for each product in the mixture. Any recommended rate of this product may be used in a tank mix.

NOTE: For forestry site preparation, make sure the tank-mix product is approved for use prior to planting the desired species. Observe planting interval restrictions. For side trimming treatments in utility rights-of-way, tank mixtures with Arsenal® 2WSL are not recommended. For side trimming treatments, it is recommended that this product be used alone as recommended, or as a tank mixture with Garlon® 4.

Product	Broadcast Rate	Use Sites
Arsenal® Applicators Concentrate**	2 to 16 fl oz/a	Forestry site preparation
Chopper®	4 to 32 oz/a	Forestry site preparation
Escort®**	½ to 3 ½ oz/a	Forestry site preparation
Oust®	1 to 4 oz/a	Forestry site preparation, Utility sites
Garlon® 3A*, Garlon® 4	1 to 4 qts	Forestry site preparation, Utility sites
Arsenal® 2WSL**	4 to 32 fl oz/a	Utility sites

Product	Spray-to-Wet Rates	Use Sites
Arsenal® Applicators Concentrate**	1/32 % to ½ % by volume	Forestry site preparation
Arsenal® 2WSL**	1/16 % to ½ % by volume	Utility sites

Product	Low Volume Directed Spray Rates	Use Sites
Arsenal® Applicators Concentrate**	1/8 % to ½ % by volume	Forestry site preparation
Arsenal® 2WSL**	1/8 % to ½ % by volume	Utility sites

*Ensure that Garlon® 3A is thoroughly mixed with water according to label directions before adding this product. Have spray mixture agitating at the time this product is added to avoid spray compatibility problems.

**Arsenal® and Escort® are not registered in the state of California.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or tough-to-control woody brush and trees, use the higher recommended rates.

AD112907

Manufactured for
Albaugh Inc.
Ankeny, IA 50021