



SUPPLEMENTAL LABEL

CLETHODIM 2E

READ THE ENTIRE LABEL FOR CLETHODIM 2E BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

When using CLETHODIM 2E as permitted according to this supplemental labeling, read and follow all applicable directions, restrictions, and precautions on the label booklet provided with the pesticide container and on this supplemental labeling. This supplemental labeling must be in the possession of the user at the time of pesticide application.

EPA Reg. No. 42750-72

AD031208

USE ON SUCCULENT BEANS

KEEP OUT OF REACH OF CHILDREN

CAUTION

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)**

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

- Do not apply CLETHODIM 2E by chemigation in the states of Idaho, Montana, Oregon and Washington
- Do not apply more than 32 fl. oz. of CLETHODIM 2E (0.50 lb. ai) per acre per season. Application on Long Island, New York, is restricted to no more than 16 fl. oz. of CLETHODIM 2E (0.25 lb. ai) per acre per season.
- Do not apply more than 8 fl. oz./A of CLETHODIM 2E per application to the following crops: bean (succulent)
- Exceeding these recommendations may result in unacceptable crop injury.

CROP SPECIFIC RESTRICTIONS AND LIMITATIONS FOR CLETHODIM 2E

Crops ⁽¹⁾	Minimum Time From Application to Harvest (PHI)	Use Rates Per Acre	Crop Oil Concentrate Rates Per Acre ⁽²⁾	Special Use Instructions
Bean, Succulent including: Bean (Phaseolus spp.) Broad Bean (succulent) Lima Bean (green) Bean (vigna spp.) Black-eyed Pea Cowpea Southern Pea	21 days	6-8 fl. oz.	1 qt. by ground or 1% v/v (but not less than 1 pt./A) by air. ⁽³⁾	Refer to appropriate Table for reduced rate recommendations for the control of small annual grasses. Do not apply more than one (1) application per acre per season. The addition of AMS has shown improved grass control for difficult to control species including: quackgrass, rhizome Johnsongrass, red rice, wild oats, volunteer cereals and volunteer corn.

N/A = Not Applicable

- (1) CLETHODIM 2E is not recommended for use on vegetable crops being grown for seed production unless specific use directions are provided.
- (2) Acceptable crop oil concentrates would be those which contain a minimum of 80% oils and 15% emulsifier. A crop oil concentrate must contain either a petroleum or vegetable oil base and must meet all the following criteria: be non-phytotoxic, contain only EPA-exempt ingredients, provide good mixing quality and be successful in local experience. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. See the Addition of Adjuvant and Crop Oil Concentrate section for further information.
- (3) 1 to 2 qts./A of liquid fertilizer (10-34-0, 28%N or 32%N), or an equivalent amount (2.5 to 4.0 lbs./A) of spray grade ammonium sulfate (AMS) may be added to CLETHODIM 2E applications, in addition to the recommended rate of crop oil concentrate.

Manufactured By:
 Albaugh, Inc.
 1525 NE 36th Street
 Ankeny, IA 50021