

MACHO 600

EPA Reg. No. 42750-133

This seed has been treated with the active ingredient **Imidacloprid** at the rates specified by the manufacturer.

CAUTION – TREATED SEED

KEEP OUT OF REACH OF CHILDREN

- Treated seed must not be used for or mixed with food or animal feed or processed for oil.
- Store away from feed and foodstuffs.
- Do not allow children, pets or livestock to have access to treated seeds.
- Do not use bag or container for refilling with foodstuff.

PRECAUTIONARY INFORMATION FOR TREATED SEED

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear Long-sleeved shirt and long pants, socks and shoes, protective eyewear and chemical-resistant gloves when handling. Remove contaminated clothing and wash before reuse.

IN CASE OF AN EMERGENCY INVOLVING THIS PRODUCT
CALL: CHEMTREC: (800) 424-9300

Albaugh, Inc.
Ankeny, IA 50021

ENVIRONMENTAL HAZARDS

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

USE RESTRICTIONS

ROTATIONAL CROPS - The following plant-back intervals should be observed:

- IMMEDIATE - Any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient including Artichoke, Barley, Brassica (cole) leafy vegetable, Canola, Cilantro, Corn, Field, Corn, Sweet, Cotton, Cranberry, Cucurbits, Eggplant, Ground cherry, Leaf petiole vegetable, Leafy vegetable, Legume vegetable (succulent or dried, except soybean), Mustard Seed, Oats, Okra, Pepinos, Pepper, Popcorn, Potato, Rapeseed, Rye, Sorghum, Soybean, Strawberry, Sugarbeet, Tomatillo, Tomato, Triticale, Vegetable Root & Tuber, Watercress, Wheat
- 30-DAY PLANT-BACK - Cereals, including: buckwheat, millet, rice, safflower
- 12-MONTH PLANT-BACK - Other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient has been established.

Do not graze or feed livestock on treated areas for 45 days after planting.