

SUPPLEMENTAL LABELING

All applicable directions, restrictions, precautions and Warranty Limitations on the EPA-registered label are to be followed.

Albaugh RANGE STAR™

EPA Reg. No. 42750-55

BETWEEN CROP APPLICATIONS, CONSERVATION RESERVE PROGRAMS, GENERAL FARMSTEAD AND FALLOW SYSTEMS

KEEP OUT OF REACH OF CHILDREN

DANGER/PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FOR CHEMICAL SPILL, LEAK, FIRE, OR EXPOSURE, CALL CHEMTREC (800) 424-9300

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Important

Before using this product, read and carefully observe all applicable directions, restrictions, and precautionary statements on the EPA-registered container label and the Range Star™ Directions for Use booklet affixed to the container.

These uses are considered Food/Feed Crops when harvested, grazed or foraged. Refer to the **ADDITIVES** section for information on adjuvant restrictions and the **NON-FOOD/FEED USE (LAND NOT HARVESTED, GRAZED OR FORAGED) – SPECIFIC INFORMATION** section for specific use directions.

NON-FOOD/FEED USE (LAND NOT HARVESTED, GRAZED OR FORAGED) – SPECIFIC INFORMATION

Between Crop Applications

PREPLANT DIRECTIONS (POSTHARVEST, FALLOW, CROP STUBBLE, SET-ASIDE) FOR BROADLEAF WEED CONTROL:

Range Star™ can be applied either postharvest in the fall, spring, or summer during the fallow period or to crop stubble/set-aside acres. Apply Range Star™ as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (postharvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer.

Refer to the **Crop Rotational Restrictions** and **General Restrictions and Limitations** for the recommended interval between application and planting to prevent crop injury.

Rates and Timings:

Apply 0.5-6 pints of Range Star™ per acre. Refer to the **ANNUAL WEEDS** and **BIENNIAL AND PERENNIAL WEEDS Application Rate and Timing** tables to determine the use rates for specific targeted weed species. Retreatments may be made as needed; however, do not exceed a total of 8 pints of Range Star™ per treated acre during a growing season. For best performance, apply Range Star™ when annual weeds are less than 6" tall, when biennial weeds are in the rosette stage and to perennial weed regrowth in late summer or fall following a mowing or tillage treatment. The most effective control of upright perennial broadleaf weeds such as Canada thistle and Jerusalem artichoke occurs if Range

Star™ is applied when the majority of weeds have at least 4-6" of regrowth or for weeds such as field bindweed and hedge bindweed that are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds that develop from seed or underground plant parts such as rhizomes or bulblets, after the effective period for Range Star™. For seedling control, a follow-up program or other cultural practices could be instituted.

Between Crop Tank Mixes

Apply 0.5-2 pints of Range Star™ per acre in tank mixes with one or more of the following herbicides for control of annual weeds, or 2-8 pints of Range Star™ per acre for control of biennial and perennial weeds:

Aim™	Glyphosate
Ally®	Gramoxone® Extra
Amber®	Kerb™
Atrazine	Landmaster® BW
Curtail™	Paramount®
Cyclone®	Sencor®
Distinct®	Tordon™ 22K
Fallow Star™	Touchdown®
Fallowmaster®	2,4-D
Finesse®	

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION.

Range Star™ is a trademark of Albaugh, Inc.

Albaugh, Inc.
121 NE 18th Street
Ankeny, IA 50021