



SUPPLEMENTAL LABELING

TEBUSTAR 3.6 FL

42750-99

READ THE ENTIRE LABEL FOR TebuStar 3.6L BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL. THIS SUPPLEMENTAL LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE DIRECTIONS FOR SPECIFIC CROPS: (Not For Use in California)

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Asparagus	Rust (Puccinia spp.)	4 to 6 fl. oz per acre
<p>Notes: Apply TEBUSTAR 3.6FL as a foliar spray to the developing ferns after harvest of spears is completed. Apply at the earliest sign of rust pustules or when weather conditions are conducive for rust development. Apply 4 to 6 fl oz of TEBUSTAR 3.6FL per acre (0.11 lb ai - 0.17 lb ai per acre) in alternation with another effective fungicide. Under conditions of severe rust pressure, use the higher rate. Repeat applications on a 14-day interval as necessary to maintain control of rust. Do not apply to harvestable spears. Do not apply within 100 days harvest in California and 180 days in all other states. Do not make more than three foliar applications per season (18 fl oz/acre or 0.51 lb ai/acre).</p> <p>General Comments: Applications may be made using ground or aerial application equipment. A 50 foot spray drift buffer zone is required for all aerial applications. For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL is a sterol demethylation inhibitor (DMI) fungicide (Group 3). Alternating TEBUSTAR 3.6FL with other DMI fungicides may lead to resistance.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Beans (fresh & dry except succulent shelled)	Rust (Uromyces appendiculatus)	4 to 6 fl. oz per acre
<p>Notes: Apply TEBUSTAR 3.6FL in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 14-day intervals, or as necessary to maintain control. Beans, fresh: TEBUSTAR 3.6FL may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of TEBUSTAR 3.6FL per acre per crop season. Beans, dry: TEBUSTAR 3.6FL may be applied up to 14 days before harvest. Do not apply more than 12 fl. oz. of TEBUSTAR 3.6FL per acre per crop season.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on bean foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p>		

Restricted-entry interval (REI) = 12 hours.

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Corn (sweet corn, field corn, field corn grown for seed, and popcorn)	Rust (<i>Puccinia</i> spp.) Northern leaf blight (<i>Helminthosporium turcicum</i>) Southern leaf blight (<i>Helminthosporium maydis</i>) Northern leaf spot (<i>Helminthosporium carbonum</i>) Gray leaf spot (<i>Cercospora zeae-maydis</i>)	4 to 6 fl. oz per acre
Notes: Apply TEBUSTAR 3.6FL in a protective spray schedule or when weather conditions are favorable for disease development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. A maximum of 24 fl. oz. (1.5 pint) of TEBUSTAR 3.6FL may be applied per acre per crop season. Sweet corn: TEBUSTAR 3.6FL may be applied up to 7 days before harvest of ears or forage, and 49 days before the harvest of fodder. Field, seed or popcorn: TEBUSTAR 3.6FL may be applied up to 21 days before the harvest of forage, and 36 days before the harvest of grain or fodder.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on corn foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) for sweet corn = 19 hours. Restricted-entry interval (REI) for all corn except sweet corn = 12 hours.		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Cotton	Southwestern cotton rust (<i>Puccinia cacabata</i>)	6 to 8 fl. oz per acre
Notes: Apply TEBUSTAR 3.6FL in a protective spray schedule or when weather conditions are favorable for rust development. Repeat applications at 7- to 14-day intervals, or as necessary to maintain control. TEBUSTAR 3.6FL may be applied up to 30 days before harvest. Do not apply more than 24 fl. oz. of TEBUSTAR 3.6FL per acre per crop season.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on cotton foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3). Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Cucurbit Vegetables Group Chayote Chinese waxgourd Citron melon Cucumber Gherkin Edible gourd, (includes hyotan, cucuzza, hechima and Chinese okra) Momordica spp. (includes balsam apple, balsam pear, bitter melon and Chinese cucumber) Muskmelon (includes cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon and snake melon) Pumpkin Summer squash (includes crookneck squash, scallop squash, straightneck squash, vegetable marrow and zucchini) Winter squash (includes butternet squash, calabaza, hubbard squash, acorn squash and spaghetti squash) Watermelon	Powdery mildew (Sphaerotheca fuliginea / Podosphaera xanthii) (Erysiphe cichoracearum)	4 to 6 fl. oz per acre
	Gummy stem blight – suppression (Didymella bryonae) (watermelon, squash, pumpkin, and melons only)	8 fl. oz. per acre
	Notes: Apply the specified dosage in a protective spray schedule to foliage and fruit. Repeat applications at 10- to 14-day intervals. TEBUSTAR 3.6FL may be applied up to 7 days before harvest. Do not apply more than 24 fl. oz. of TEBUSTAR 3.6FL per acre per crop season.	
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		
CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Hops	Powdery mildew (Sphaerotheca humuli/ Sphaerotheca macularis)	4 to 8 fl. oz per acre
Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 10- to 14-day intervals. TEBUSTAR 3.6FL may be applied up to 14 days before harvest. Do not apply more than 32 fl. oz. of TEBUSTAR 3.6FL per acre per crop season. Increase the spray volume and the application rate as vine growth increases during the season.		
General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).		

Restricted-entry interval (REI) = 12 hours.		
CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Green Onion Leek Spring Onion Scallion Japanese bunching onion Green shallots Green eschalots	White rot (Sclerotium cepivorum) suppression only Rust (Puccinia allii, Puccinia porri) Purple blotch (Alternaria porii)	4 to 6 fl. oz per acre
For the control of diseases make foliar applications using an interval of 10 to 14 days. Apply TEBUSTAR 3.6FL in a protective spray schedule or when weather conditons are favorable for rust development. Notes: Do not apply more than 24 fl. oz. of TEBUSTAR 3.6FL per acre per crop season. Do not apply within 7 days of harvest (PHI = 7 days).		
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		
CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Leafy Brassica Greens Broccoli raab Chinese cabbage (bok choy) Collards Kale Mizuma Mustard greens Mustard spinach Rape greens Turnip greens	Cercospora leaf spot (Cercospora brassicicola) Powdery mildew (Erysiphe cruciferarum) Alternaria leaf spot (Alternaria brassicicola)	3 to 4 fl. oz per acre
Notes: Do not apply more than 16 fl. oz. of TEBUSTAR 3.6FL per acre per crop season. Do not apply within 7 days of harvest (PHI = 7 days)		
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3). Restriction: Application to turnip greens is limited to East of the Rockies.		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Garden beet roots and tops (leaves)	Cercospora leaf spot (Cercospora beticola)	3 to 7.2 fl. oz per acre
	Notes: Make applications on a 14 day intervals. Do not apply more than 28.8 fl. oz. TEBUSTAR 3.6FL per acre season. Do not apply within 7 days of harvest (PHI = 7 days).	
General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).		
Restricted-entry interval (REI) = 12 hours.		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Lychee	Anthrachnose (Colletotrichum gloeosporioides)	4 to 6 fl. oz per acre
Notes: Begin first application of TEBUSTAR 3.6FL as panicle emerges. Spray up to 6 fl. oz. per acre every 10 days thereafter for a total of 8 sprays. Apply specific dosage in a minimum of 50 gallons of spray solution per acre by ground only. Do not apply more than 48 fl. oz. of TEBUSTAR 3.6FL per acre season. TEBUSTAR 3.6FL can be applied up to and including the day of harvest (PHI = 0 days).		
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 2 days.</p>		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Okra	Cercospora leaf spot (Cercospora spp.)	4 to 6 fl. oz per acre
Notes: Apply specified dosage of TEBUSTAR 3.6FL in a preventative spray program. Use the highest rate when disease conditions are favorable and in areas where high disease pressure is expected. Applications may be repeated at 14-day intervals in order to maintain control of the disease. Apply specified dosage as a foliar spray in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Applications may be made no closer than 3 days before harvest. Do not apply more than 24 fl. oz. of TEBUSTAR 3.6FL per acre per season.		
<p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Pecan	Brown leaf spot (Sirosporium diffusum) Downy spot (Mycosphaerella caryigena) Liver spot (Gnomonia caryae) Scab (Cladosporium caryigenum) Vein spot (Gnomonia nerviseda) Zonate leaf spot (Grovesinia pyramidalis)	4 to 8 fl. oz per acre
<p>Notes: Apply TEBUSTAR 3.6FL in a preventive spray schedule beginning at early bud break (young leaves unfolding), and continue applications at 10- to 14-day intervals through the pollination period. TEBUSTAR 3.6FL may be applied at 4 fl. oz. per acre in tank-mix with the labeled rate of Super-Tin® in cover sprays. Follow label directions for the use of Super-Tin. Do not add a surfactant to the spray solution when tank-mixing TEBUSTAR 3.6FL with Super-Tin. Apply TEBUSTAR 3.6FL in a spray volume of 15 or more gallons per acre by air or 50 or more gallons per acre by ground. Apply 7 to 8 fl. oz. per acre of TEBUSTAR 3.6FL to full-size mature trees, and 4 to 6 fl. oz. per acre of TEBUSTAR 3.6FL to smaller trees. Apply the high rate to varieties that are highly susceptible to the indicated diseases, or when severe disease conditions exist. The lowest labeled rate of a surfactant may be added to the spray solution for optimum control of the indicated diseases. Do not apply after shucks begin to split. A maximum of 32 fl. oz. of TEBUSTAR 3.6FL may be applied per acre per crop season. Do not cut cover crops in treated areas for feed or allow livestock to graze treated areas.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3). It may be applied in a tank-mix or alternated (every other spray application) with a non-DMI fungicide as a resistance management strategy.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		
CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Soybean	Rust (Phakopsora pachyrhizi) Powdery mildew (Microsphaera diffusa)	3 to 4 fl. oz per acre
<p>Use Directions: Apply TEBUSTAR 3.6FL as a broadcast foliar spray as a preventative spray or at first visible symptoms of disease. Repeat applications on a 10- to 14-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. The lowest label labeled rate of a spray surfactant must be tank-mixed with Folicur 3.6F. Folicur 3.6F should be applied in a minimum for 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.</p> <p>Restrictions: Applications may not be made within 21 days of harvest. Do not apply more than 3 applications per season. Do not apply ore than 12 fl. oz./A per use season.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Sunflower	Rust (Puccinia helianthi)	4 to 6 fl. oz per acre
<p>Notes: Apply specified dosage of TEBUSTAR 3.6FL at the earliest sign of infection (rust pustules developing) or when weather conditions are favorable for rust development. Apply higher rate to highly susceptible varieties and/or under severe disease conditions. Apply specific dosage in a minimum of 20 gallons of spray solution per acre by ground or a minimum of 5 gallons of spray solution by air. Do not apply more than 16 fl. oz. of TEBUSTAR 3.6FL per acre per crop season or within 50 days of harvest.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. Contact your state Extension Service or Bayer representative for a list of approved surfactants. TEBUSTAR 3.6FL must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Turnip (Application is limited to East of the Rockies)	Cercospora leaf spot (Cercospora brassicicola)	4 to 7.2 fl. oz per acre
<p>Notes: Apply the specified dosage in a protective spray schedule to foliage. Repeat applications at 12- to 14-days intervals. TEBUSTAR 3.6FL may be applied up to 7 days before harvest. Do not apply more than 28.8 fl. oz. of TEBUSTAR 3.6FL per acre per crop season.</p> <p>General Comments: For optimum disease control, the lowest labeled rate of a spray surfactant should be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on plant foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restriction: Application to turnip greens is limited to East of the Rockies.</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

Manufactured by:
Albaugh, Inc.
Ankeny, IA 50021

AD031909