



SUPPLEMENTAL LABELING

TEBUSTAR 3.6 FL

42750-99

READ THE ENTIRE LABEL FOR TebuStar 3.6L BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL. THIS SUPPLEMENTAL LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

KEEP OUT OF REACH OF CHILDREN

CAUTION

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

USE DIRECTIONS FOR SPECIFIC CROPS: (Not For Use in California)

CROP	DISEASE	RATE OF TEBUSTAR 3.6FL
Dry bulb onion Garlic Great-headed (elephant) garlic Welch onion Shallot	White rot (Sclerotium cepivorum)	White rot: 20.5 fl oz per acre applied in a 4 to 6 inch band over/into each furrow. May be applied by chemigation to control white rot.
	Rust (Puccinia allii, Puccinia porri)	4 to 6 fl. oz. per acre
	Purple blotch (Alternaria porii)	
<p>White rot: For the control of white rot, make one application in the furrow at the time of planting. The in-furrow application should be made at the rate of 20.5 fl. oz. TEBUSTAR 3.6FL per acre. Apply the entire per acre rate in a 4 to 6 inch band over/into each furrow. Additional control may be obtained by including two foliar applications at 4 to 6 fl oz/acre.</p> <p>Rust: For the control of rust make foliar applications at the rate of 4 to 6 fl. oz TEBUSTAR 3.6FL per acre per application. Repeat at an interval of 10 to 14 days.</p> <p>Apply TEBUSTAR 3.6FL in a protective spray schedule or when weather conditions are favorable for rust development.</p> <p>Notes: Do not apply more than 32.5 fl. oz. of TEBUSTAR 3.6FL per acre per crop season if an in-furrow treatment is made. If TEBUSTAR 3.6FL is not applied as an in-furrow treatment then do not apply more than 12 fl oz. TEBUSTAR 3.6FL per acre per season as a foliar spray. Do not apply within 7 days of harvest (PHI = 7 days).</p> <p>General Comments: For optimum results use as a preventative treatment. Begin applications as soon as crop and/or environmental conditions become favorable for disease development. The lowest labeled rate of a spray surfactant may be tank-mixed with TEBUSTAR 3.6FL. TEBUSTAR 3.6FL must have two to four hours of drying time on foliage for the active ingredient to move systemically into plant tissue before rain or irrigation occurs. After this period of time, TEBUSTAR 3.6FL will be resistant to weathering. TEBUSTAR 3.6FL is a demethylation inhibitor (DMI) fungicide (Group 3).</p> <p>Restricted-entry interval (REI) = 12 hours.</p>		

Manufactured by:
Albaugh, Inc.

AD031909

Ankeny, IA 50021