

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for the uses covered by the certified applicator's certification.

FANFARE[®] 2EC

EPA REG. NO. 66222-99

FIFRA 2(ee) RECOMMENDATION FOR CONTROL OF VARIOUS PESTS IN SOYBEANS

For applications only in Alabama, Arkansas, Florida, Georgia, Iowa, Illinois, Indiana, Kansas, Kentucky, Louisiana, Missouri, Minnesota, Mississippi, Nebraska, North Carolina, South Carolina, South Dakota, Tennessee, Wisconsin

DIRECTIONS FOR USE

This FIFRA Section 2(ee) Recommendation recommends using Fanfare 2EC on soybeans for control and/or suppression of listed pests. The user must refer to the federally approved Fanfare 2EC label and read and follow all directions for use, restrictions, and precautions. This recommendation is permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA. It is a violation of federal law to use this product in a manner inconsistent with its labeling. Users should have this recommendation in their possession at the time of use.

PREHARVEST INTERVAL: The required days to wait between the last application and harvest are given in () after the crop name.

SOYBEAN (18)

PEST	DOSAGE		COMMENTS
	LB AI/A	FL OZ/A	
Soybean Aphid Alfalfa Caterpillar, Aphids, Aster Leafhopper, Bean Leaf Beetle, Beet Armyworm*, Cloverworm, Corn Earworm, Corn Rootworm Adult, Cucumber Beetle, Cutworms, European Corn Borer, Fall Armyworm, Flea Beetle, Grasshoppers, Imported Cabbageworm, Japanese Beetle Adult, Leafhoppers, Leafminers, Loopers, Mexican Bean Beetle Adult, Pea Leaf Weevil, Plant Bug, Saltmarsh Caterpillar, Sap Beetle, Southern Armyworm, Stink Bugs, Tarnished Plant Bug, Thrips, Tobacco Budworm*, Webworms, Western Bean Cutworm, Whitefly, Yellowstriped Armyworm	0.033-0.10	2.1-6.4	Ground application: Apply in water in a minimum of 10 gallons per acre. Air application: Apply in water in a minimum of 2 gallons per acre. *Pyrethroid resistance is common for Beet Armyworm and Tobacco Budworm. Consult with state or local extension service representatives to determine if resistance pest populations are present in your area. If resistance has been detected in your area, refer to the RESISTANCE MANAGEMENT statement found on the label attached to the container for further guidance.
Two spotted Spider Mite Lygus Species Whitefly	0.08-0.10	5.12-6.4	

- To maintain a proper spray interval, do not make applications less than 30 days apart.
- Do not apply more than 0.3 lb. active ingredient (12.8 ounces formulated) per acre per season.

REV 072810



4515 Falls of Neuse Road, Suite 300
Raleigh, NC 27609