

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON



EPA REG. NO. 10163-251 / SLN NO. WA-010026

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- This label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions and precautions on the EPA-registered label.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants in farms, forests, nurseries, or greenhouses.

REENTRY: Keep children, pets and unprotected persons out of treated areas until sprays have dried.

DIRECTIONS FOR USE

Hexygon DF is predominantly active against eggs and immature mites and should be applied prior to adult mite buildup. Control is achieved from either direct contact with the spray or from contact with treated plant surfaces. Good coverage is essential for optimum performance with Hexygon DF. Use sufficient volume to obtain thorough, uniform coverage. Spray all foliage surfaces including the underside of leaves for complete coverage.

CROP	PEST	RATE	COMMENTS
Residential Pome Fruit Trees, Stone Fruit Trees And Nut Trees	European Red Mite, Twospotted Spider Mite, Willamette Mite	1 -2 oz. per 100 gals. applied dilute OR 3 - 6 oz. per acre applied concentrate OR 0.07-0.14 oz. per 1000 square feet	Apply lower rates when mite infestation levels are low and higher rates on moderate to high mite infestation levels. Make only one application per year. This product is intended for use in home plantings by licensed commercial pesticide applicators.

- Do not apply within 28 days of harvest.
- For use under this SLN label, do not apply this product through any type of irrigation system.

This pesticide is toxic to fish and aquatic invertebrates. Hexygon DF should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

Plastic containers - WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

This label for Hexygon DF expires and must not be distributed or used in accordance with this SLN registration after December 31, 2011.

®Hexygon Ovicide/Miticide is a trademark of Gowan Company.
24(c): Registrant
Gowan Company
P.O. Box 6659
Yuma, AZ 85364

Approved: 10-18-06