

FIFRA §2(ee) RECOMMENDATION
FOR USE ONLY IN CALIFORNIA

RESTRICTED USE PESTICIDE

Due to Toxicity to Fish, Birds, and Aquatic Organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

Mesuro[®]l 75-W

Insecticide - Miticide - Molluscicide

EPA Reg. No. 10163-231

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.

For control of Glassy-Winged Sharpshooter and Citrus Thrips

General Precautions and Restrictions: Do not use on crops for food or forage. Use only according to label directions. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR APPLICATION

CROP	PEST	LBS./ 100 GALS.	COMMENTS
Ornamental Plants & Nonbearing Trees & Vines Growing in Nurseries and Greenhouses and Mature Ornamental Plantings	Aphids*, Mites*, Western Flower Thrips, Glassy-winged Sharpshooter, Citrus Thrips	½ - 1	Foliar Application: To control pests on foliage, flowers, and other ground plant parts, spray plants and foliage until thoroughly wet. Do not make more than 2 applications per year per crop. Applications must be at least 10 days apart.
	Snails, Slugs	2	

- Use extreme caution when applying in conjunction with foliar fertilizer applications.
- Before treating a large number of ornamental plants with Mesuro[®]l alone or as a tank mixture with any other material, make a test application on a few plants and observe for 7-10 days prior to treating large areas to reduce the possibility of plant injury.
- Do not make more than 2 applications per year per crop. Applications must be at least 10 days apart.

*In some cases these pests may have developed resistance. Mesuro[®]l used alone may not provide satisfactory control in these areas. Consult your local Agricultural Advisor or Cooperative Extension Service for recommendations.

Performance may be reduced when the spray solution has a pH greater than 7. The pH of the spray solution must be corrected by the addition of a suitable buffering or acidifying agent for optimum activity.

For more information, contact your local Gowan Company sales representative or the Gowan office.