

**Supplemental not yet approved in California**

# Onager<sup>®</sup> Miticide

**EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY**

<b>ACTIVE INGREDIENT:</b> Hexythiazox	<b>% By Wt.</b>
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide .....	11.8%
<b>OTHER INGREDIENTS</b> .....	<b>88.2%</b>
<b>TOTAL</b>	<b>100.00%</b>

Onager contains 1.0 lb. active ingredient per gallon  
Contains Petroleum Distillate

## KEEP OUT OF REACH OF CHILDREN CAUTION

### DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

### PREHARVEST INTERVAL

The required days between the last application and harvest are given in ( ) after each crop name.

### APPLICATION DIRECTIONS

CROP	PEST	RATE OZS./ACRE	COMMENTS
<b>CORN, FIELD (45)</b> (Western US <i>Only</i> *)	Banks grass mite, Two spotted spider mite, Pacific spider mite, Strawberry spider mite	10 - 24	Apply Onager at the first sign of mites before populations begin to build. <b>Ground Application:</b> Apply the recommended rate of Onager by ground equipment at 15 – 20 GPA. Use a spray boom equipped to position one 16-inch boom drop extension per row, with two nozzles directed to spray up and under the crop canopy - one to the right and one to the left. <b>Air Application:</b> Apply the recommended rate of Onager by air in a minimum of 5 GPA. Applications made by air to dense foliage may not provide adequate coverage of lower leaf surfaces for sufficient control. Use of higher labeled rates may be necessary. <b>Chemigation:</b> see details under Chemigation Systems on Section 3 label.
	<ul style="list-style-type: none"> <li>• Do not apply more than a total of 24 oz. formulated product per acre per year.</li> <li>• Do not make more than one application per year.</li> <li>• Do not harvest field corn forage, stover, and grain for 45 days after application.</li> <li>• Application after the VT crop growth stage is prohibited.</li> </ul> <p>*May only be applied in the Western US - west of the line defined as follows: TX: W of Rt. 283 and NW of Rt. 377 OK: W or Rt. 281/183 KS, NE, SD, ND: W of Rt. 281</p>		

### RESTRICTIONS ON ALL CROPS

- Do not graze or feed livestock on cover crops growing in treated areas.
- Do not use water from treated fields for irrigating other crops.
- Onager is an emulsifiable concentrate to be diluted with water for application in commercial plantings only.
- Apply Onager prior to adult mite build up. Onager will not control adult spider mites. Use higher rates on moderate to high mite infestations or for larger plants with a dense canopy. If adult mites are present in medium to high populations, better results may be obtained using Onager in combination with a registered contact adulticide. The use of less than label rates is not recommended with Onager.
- Do not make more than one application of Onager per year.
- Do not plant rotational crops other than those on this label within 120 days of this application.

### NET CONTENTS \_\_\_\_\_



EPA Reg. No. 10163-277  
EPA Est. No. 67545-AZ-01

Produced For:  
Gowan Company  
P.O. Box 5569  
Yuma, AZ 85366