

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Dithane[®] F-45 Rainshield[®]

EPA Reg No. 62719-396

Disease Control in Various Crops

Directions for Use

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Dithane[®] F-45 Flowable Mancozeb Agricultural Fungicide fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Dithane F-45 Flowable Mancozeb Agricultural Fungicide according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Dithane F-45 Flowable Mancozeb Agricultural Fungicide.

Crops	Disease Controlled	Rate per Application (qt product/acre)	Remarks (Also Refer to Directions for Use)	Restrictions
atemoya cherimoya custard apple sugar apple sweetsop	anthracnose	1.6 - 1.8	Begin applications at flowering and continue at a 7-day re-treatment interval. Applications made with aerial equipment must be made in a minimum spray volume of 10 gallons per acre.	<ul style="list-style-type: none"> • Do not apply more than 26.25 quarts of this product (26.25 lb ai) per acre per year. • Do not make more than 14 applications per year. • Applications may be made up to the day of harvest.
ginseng	alternaria blight	1.5	Begin applications when disease first threatens and repeat every 7 to 10 days as needed. In Wisconsin, apply with ground equipment and a minimum of 80 gallons of water per acre.	<ul style="list-style-type: none"> • Do not apply more than 18 quarts of this product (18 lb ai) per acre per year. • Do not make more than 12 applications per year. • Preharvest Interval: Do not apply within 30 days of harvest.
cucurbit crop group chayote Chinese waxgourd citron melon cucumber gherkin gourd, edible <i>Momordica</i> spp. muskmelon pumpkin	alternaria leaf spot anthracnose cercospora leaf spot downy mildew gummy stem blight scab	1.6 - 2.4	Begin applications when the plants are in the two-leaf stage and repeat at 7- to 10-day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. Some cantaloupe varieties (i.e., harvest queen, gold star, super star, sweet	<ul style="list-style-type: none"> • Do not apply more than 19.2 quarts of this product (19.2 lb ai) per acre per year. • Do not make more than 8 applications per year. • Preharvest Interval: Do not apply within 5 days of harvest.

Crops	Disease Controlled	Rate per Application (qt product/acre)	Remarks (Also Refer to Directions for Use)	Restrictions
squash, summer squash, winter watermelon			and early, and satcoy) are sensitive to this product. Applications made with aerial equipment must be made in a minimum spray volume of 2 gallons per acre. Consult State Cooperative Extension Service Specialist prior to use.	
canistel mamey sapote mango sapodilla star apple (caimito) white sapote	anthracnose black spot (cercospora) phytophthora fruit rot	1.6 - 2	Begin applications at flowering and continue at 14- to 21-day intervals. Direct spray to crown and blossom area. Use 20 to 100 gallons of water per acre.	<ul style="list-style-type: none"> • Do not apply more than 28 quarts of this product (28 lb ai) per acre per year. • Do not make more than 14 applications per year. • Applications may be made up to the day of harvest.

®Trademark of Dow AgroSciences LLC

R821-005

EPA accepted: 09/30/10

Initial printing.