

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Restricted Use Pesticide

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

Kerb[®] SC

EPA Reg. No. 62719-578

Preemergence and Postemergence Control of Certain Winter Annual and Perennial Grasses in Blackberry, Boysenberry, Raspberry

(For Use in Idaho)

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Kerb[®] SC herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Kerb SC according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Kerb SC.

Directions for Use

General Information

Kerb SC is a selective herbicide recommended for fall and winter applications to established blackberries, boysenberries and raspberries for both preemergence and postemergence control of certain winter annual and perennial grasses.

Dosage

Kerb SC may be applied at the rate of 2.5 to 7.0 pints of product (1 to 3 lb active ingredient) per acre broadcast application. The rate will depend on the weed species present and the soil texture of the site being treated. Follow the weed control instructions listed in the chart below:

Pints of Kerb SC SC Per Broadcast Acre ¹		
Weeds Controlled	Dependable Rainfall or Overhead Irrigation ²	Comments
bluegrass, annual	2.5 – 5.0	Use low rates on light
quackgrass	5.0 – 7.0	to medium soils and high
ryegrass, perennial	5.0 – 7.0	rates on heavy soils

¹ Dosage rates specified are in pounds of Kerb SC per acre broadcast application. Reduce rates accordingly for banded applications.

² For effective weed control, rainfall or overhead irrigation is essential following the application of Kerb SC.

Crop Tolerance

Established cane fruit are tolerant to specified rates of Kerb SC SC. Newly transplanted blackberries, boysenberries and raspberries should be well rooted and transplanted for at least 3 months prior to the application of Kerb SC.

Timing and Application

Apply Kerb SC only during the fall or winter months. For optimum results, apply Kerb SC SC during November or December. Do not make applications when the ground is frozen. Mix the specified amount of Kerb SC in clean water and apply uniformly with a low pressure ground sprayer in 20 to 50 gallons of water per acre.

Blackberry/Boysenberry/Raspberry - Specific Use Restrictions

- Do not apply more than 3 lb/acre active ingredient (7.0 pt/acre of Kerb SC) or make more than one application of Kerb SC per season.

®Trademark of Dow AgroSciences LLC

R361-010
EPA accepted: 10/18/07
Initial printing.