

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Durango[®] DMA[®]

EPA Reg. No. 62719-556

Control and Management of Glyphosate-Resistant Horseweed (*Marestail, Conyza canadensis*)

(Not for Use or Distribution Within the State of California)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Read the Product Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, Application Equipment and Application Methods, and Roundup Ready Crops on the label attached to this product's container.

Corn - Preplant, Preemergence, At-Planting

Apply this product in a preplant application to control and manage glyphosate-resistant horseweed (*marestail, Conyza canadensis*). Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. Use 24 fl oz of this product per acre in a tank mix with 2,4-D (0.5 lb active ingredient per acre) before horseweed is more than 6 inches in height. Dicamba may be used in a tank mix with this product. Include atrazine (1 to 2 lb active ingredient per acre) in the tank mix to provide residual control.

Corn Hybrids with Roundup Ready[®] 2 Technology - In-Crop

Apply this product in a post-directed application to control and manage glyphosate-resistant horseweed (*marestail, Conyza canadensis*). Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. Use 24 fl oz of this product per acre in a tank mix with 8 to 16 fl oz of Clarity per acre or 0.5 to 1 lb 2,4-D per acre between corn emergence and the 5-leaf state of growth, or when the corn is approximately 8 inches tall. Dicamba may also be included in a tank mix with this product.

Cotton - Preplant

Apply this product in a preplant application to control and manage glyphosate-resistant horseweed (*marestail, Conyza canadensis*). Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. Use 24 fl oz of this product per acre in a tank mix with 8 fl oz of Clarity per acre. Make this application 21 to 35 days before planting cotton and before horseweed reaches 6 inches in height. Observe a minimum of 21 days between the Clarity application and planting cotton during which there is at least 1 inch of cumulative rainfall in order to avoid crop injury. 2,4-D may also be included in a tank mix with this product.

Cotton with Roundup Ready[®] Gene and Roundup Ready[®] Flex Gene- Post-Directed

Apply this product in a post-directed application to control and manage glyphosate-resistant horseweed (*marestail, Conyza canadensis*). Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. Management of early season weed competition

and the development of a crop height differential between cotton and horseweed occurs by a combination of preplant burndown and postemergence over the top and/or directed applications of this product. The development of a height differential is necessary to successfully make post-directed treatments. Make in-crop post-directed applications of MSMA (2 lb active ingredient per acre) tank mixed with diuron (0.5 to 0.75 lb active ingredient per acre) when the temperature is 80°F or hotter.

Grapes

Use this product to control and manage glyphosate-resistant horseweed (marestail, *Conyza canadensis*). Apply 48 fl oz of this product per acre plus 2,4-D (18 fl oz of Dri-Clean per acre) before marestail is more than 6 inches in height. For best results, use a carrier volume of 15 gallons per acre. A residual herbicide such as diuron may provide additional preemergence control. Further local restrictions may apply.

Soybean - Preplant

Apply this product in a preplant application to control and manage glyphosate-resistant horseweed (marestail, *Conyza canadensis*). Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. For best results, control horseweed prior to planting. Use 24 fl oz of this product per acre in a tank mix with 2,4-D (0.5 lb active ingredient per acre) before horseweed is more than 6 inches in height.

Soybean with Roundup Ready® Gene and Roundup Ready® 2 Gene- In-Crop

Apply this product as a salvage treatment to glyphosate-resistant horseweed (marestail, *Conyza canadensis*) that was not controlled by a preplant application. Apply in 10 to 20 gallons of water per acre for ground applications or in 3 to 15 gallons of water per acre for aerial applications. Horseweed must be 6 inches or less in height. Apply 24 fl oz of this product per acre in a tank mix with 0.3 oz of FirstRate per acre. Apply between full emergence of the first trifoliate leaf and 50 percent flowering stage of soybeans.

Tree Fruits (Pome Fruit and Stone Fruit)

Use this product to control and manage glyphosate-resistant horseweed (marestail, *Conyza canadensis*) in all tree fruits listed in the Tree Fruits section on the label attached to this product's container. Apply 3 pints of this product per acre plus 2,4-D (18 fl oz of Dri-Clean per acre or 2 pints of Orchard Master CA per acre) before marestail is more than 6 inches in height. For best results, use a carrier volume of 15 gallons per acre.

Tree Nuts

Use this product to control and manage glyphosate-resistant horseweed (marestail, *Conyza canadensis*) in all tree nuts listed in the Tree Nuts section on the label attached to this product's container. Apply 3 pints of this product per acre plus 2,4-D (18 fl oz of Dri-Clean per acre or 2 pints of Orchard Master CA per acre) before marestail is more than 6 inches in height. For best results, use a carrier volume of 15 gallons per acre.

®Trademark of Dow AgroSciences LLC
Roundup Ready® is a registered trademark of Monsanto Company

R353-045
EPA accepted 11/26/10
Replaces R353-025.