

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Durango[®] DMA[®]

EPA Reg. No. 62719-556

Weed Control in Cotton with Roundup Ready[®] Gene in Arizona

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for this product before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of this product according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use

Read the General Information, Herbicide Resistance Management, Spray Drift Management, Mixing Directions, Application Equipment and Application Methods, and Roundup Ready Crops on the label attached to this product's container.

See the Roundup Ready Crops section of the product label for general precautionary instructions for use in Roundup Ready Crops.

Types of Applications: Preplant, at-planting, preemergence, over the top, selective equipment, preharvest

Maximum Allowable Application Rates

Application Type	Rate (quart/acre)
preplant preemergence at-planting	3.75
maximum preharvest	1.5
total all in-crop applications from ground cracking to layby	2.75
total in-crop over the top from ground cracking to 4-leaf stage	2.25
total in-crop applications using selective equipment through layby	1.5
combined total for all applications	6

Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 7 days before harvest.
- Do not make more than two over the top broadcast applications from crop emergence through the 4-leaf (node) development stage.
- Do not make more than two post-directed applications from the 5-leaf stage through layby.
- There must be 10 days between sequential in-crop over the top or post-directed applications of this product. Cotton must also have at least two nodes of incremental growth between applications.
- Do not apply more than the maximum label rates of this product. Doing so will result in boll loss, delayed maturity and/or yield loss and are the sole responsibility of the grower.

- The maximum use rates on this label apply to the rates of this product combined with the use of all other glyphosate-containing products, applied separately or as a tank mix. Calculate the application rates and ensure that the total use of this and other glyphosate-containing products does not exceed the specified maximum use rate.

Preplant, Preemergence, and At-Planting

Apply this product before, during, or after planting Roundup Ready cotton.

Over the Top

Apply up to 24 fl oz of this product per application using aerial or ground application equipment. Make this postemergence application to Roundup Ready cotton from the ground cracking stage until the 4-leaf (node) development stage (until the fifth true leaf reaches the size of a quarter).

Precautions and Restrictions:

- Do not apply more than 24 fl oz of this product per acre in any single over the top broadcast application.
- Do not apply more than 2.25 quarts of this product per acre for combined over the top applications between ground cracking until the 4-leaf (node) stage.

Selective Equipment

Apply this product using precision post-directed or hooded sprayers to Roundup Ready cotton through layby using up to 24 fl oz per acre per application. Up to 1.5 quarts of this product per acre may be used in sequential in-crop applications using selective equipment. Use post-directed equipment that directs the spray to the base of the cotton plants. To avoid contact of the herbicide spray with leaves of the cotton plant to the maximum extent possible, maintain a low spray pressure (less than 30 lb psi) and place the nozzles in a low position. This directs a horizontal spray pattern under the leaves of the cotton plant and onto the weeds in the row. For best results, apply while weeds are small and less than 3 inches in height.

Precautions and Restrictions:

- Do not apply more than a total of 2.75 quarts of this product per acre as a combined in-crop over the top plus selective equipment applications

Salvage Treatment: Apply up to 36 fl oz of this product per acre from the ground cracking stage through layby when weeds threaten to cause the loss of the crop. Make the application either as an over the top or as a post-directed treatment sprayed higher on the cotton plants and over the weeds. **Note:** Crop tolerance of Roundup Ready cotton has not fully been tested at this application rate. A salvage treatment will result in significant boll loss, delayed maturity and/or yield loss and are the sole responsibility of the grower. Do not apply more than two salvage treatment per growing season.

Precautions and Restrictions:

- Do not apply more than a total of 2.75 quarts per acre of this product in a combined in-crop and over the top plus selective equipment applications.

Preharvest

This product may be applied for preharvest annual and perennial weed control as a broadcast application after 20 percent of boll crack. This product does not enhance the performance of harvest aids when applied to Roundup Ready cotton.

Precautions and Restrictions:

- **Preharvest Interval:** Do not apply within 7 days of harvest.
- Do not apply more than 1.5 quarts of this product using either aerial or ground application equipment.
- Do not apply this product preharvest to Roundup Ready cotton grown for seed.

Attention: Use this product as an over the top or directed application onto cotton that is designated as Roundup Ready. If cotton varieties not designated as Roundup Ready are sprayed with this product, severe injury or death will result. Avoid contact of herbicide with foliage, green stems, or fruit of crops, or any desirable plants and trees other than crops with a Roundup Ready gene as severe injury or destruction will result.

®Trademark of Dow AgroSciences LLC
Roundup Ready® is a registered trademark of Monsanto Company
R353-028
EPA accepted 11/26/10
Initial printing