
SUPPLEMENTAL LABELING

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1226A-S1 0709

Alto® 100SL

Fungicide

Broad spectrum fungicide for control of plant diseases

Active Ingredient:	
Cyproconazole (CAS No. 94361-06-5).....	8.9%
Other Ingredients:	91.1%
Total:	100.0%

Alto 100SL is a soluble concentrate formulation which contains 0.83 lb. a.i. cyproconazole per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1226

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Alto 100SL as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Wheat Triticale	Early season suppression of: Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nordorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>)	1.5–5.5	Apply Alto 100SL in the spring at approximately Feekes Stage 5. Applications may be made no closer than a 14-day interval. Alto 100SL may be applied by ground, air or chemigation.
	Control of Leaf Diseases: Rust (<i>Puccinia</i> spp.) Powdery mildew (<i>Blumeria</i> spp., <i>Erysiphe</i> spp.) Leaf Blight (<i>Septoria tritici</i>) Glume Blotch (<i>Stagonospora nordorum</i>) Tan spot (<i>Pyrenophora tritici-repentis</i>) Helminthosporium leaf blight (<i>Drechslera tritici-repentis</i>) Spot Blotch (<i>Bipolaris sorokiniana</i>)	3.0–5.5	For best results, apply Alto 100SL between Feekes Stage 8 and 10.5.1. Protecting the flag leaf is important for maximizing the potential yield. If disease pressure is low, 3 fl. oz./acre may be applied. Applications may be made no closer than a 14-day interval. Alto 100SL may be applied up to 30 days prior to harvest. Alto 100SL may be applied by ground, air or chemigation.
	Foot Rot (<i>Pseudocercospora</i> spp.)	3.0–5.5	Apply Alto 100SL plus half rates of other EPA registered fungicides such as Topsin® M. Apply at tillering, but before elongation has occurred.

Specific Use Restrictions:

- (1) Do not apply more than 5.5 fl. oz./acre of Alto 100SL per season.
- (2) Do not harvest forage or hay within 21 days of an application.
- (3) Do not harvest wheat within 30 days of application.
- (4) Do not apply more than 0.036 lb. a.i./A/year cyproconazole containing products.

Alto® and the Syngenta logo are trademarks of a Syngenta Group Company

©2010 Syngenta