



ATRA

Section 24(c) Special Local Need Label

**RESTRICTED USE PESTICIDE
(GROUND AND SURFACE WATER CONCERNS)**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER.

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
OKLAHOMA**

AATREX® 4L

**EPA Reg. No. 100-497
EPA SLN No. OK-910003**

For Control Of Pigweed In Grain And Forage Sorghum.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including Worker Protection Standards, and precautions on the EPA registered AAtrex 4L label.

This labeling must be in possession of the user at the time of application.

AAtrex 4L for Control of Pigweeds in Grain and Forage Sorghum

AAtrex 4L applied by ground equipment to the soil surface as a broadcast or band treatment, may be used on medium- to fine-textured soils to control pigweeds in grain or forage sorghum or on ground to be planted to grain or forage sorghum. When using a band treatment, reduce the amount of AAtrex 4L per acre that would be broadcast in proportion to the band width of the area to be treated.

Broadcast 1-1.2 quarts of AAtrex 4L preemergence to sorghum and pigweeds in areas except northeast Oklahoma. In northeast Oklahoma, follow the use directions on the federal AAtrex 4L label. Use the higher rate in the rate range on finer-textured soils.

Note: **DO NOT** apply this product by aerial application.

Precautions: To reduce the potential for injury to sorghum, 1) **DO NOT** use on sand, loamy sand, and sandy loam soils. 2) **DO NOT** use on soils containing less than 1% organic matter. 3) **DO NOT** use on soils with pH above 7.5. 4) **DO NOT** use on soils with exposed calcareous subsoil, or on caliche or rocky outcroppings. 5) **DO NOT** use on sorghum either planted in or to be planted in furrows.

Notes: 1) Applications of AAtrex to grain sorghum on highly alkaline soils may result in crop injury. 2) Moisture is essential following application to activate AAtrex and result in control of pigweed. If dry conditions follow the AAtrex applications and pigweed plants emerge, cultivation is recommended to improve control. 3) Injury to sorghum may occur if the amount of rainfall soon after application of AAtrex is sufficient to leach it into the germinating seed zone. 4) Injury to sorghum may occur if AAtrex is applied preemergence following an at-planting systemic insecticide used in-furrow. 5) Injury to sensitive rotational crops, other than corn or sorghum, may occur the following year if prolonged periods of extremely dry weather follow the use of AAtrex.

Rotational Crops: 1) **DO NOT** rotate to any crop except corn or sorghum following treatment with AAtrex until the following year or illegal residues may result. 2) **DO NOT** plant sugar beets, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following application, or injury may occur.

Chemigation: **DO NOT** apply this product through any type of irrigation system.

AAtrex® trademark of Syngenta

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: OK-910003

Label Code: OK0497023BA1103

Word: OKAAAtrex4LGrainSorghumSLNLbl9-17-03