



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA

QUADRIS® FLOWABLE FUNGICIDE

EPA Reg. No. 100-1098

EPA SLN NO. MT-06-0006

For control of Alternaria leaf spot (*Alternaria spp.*) and Downy mildew (*Plasmopora halstedii*, *Plasmopora helianthi*) on Safflower

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Quadris Flowable Fungicide label (EPA Reg. No. 100-1098).

This label must be in the possession of the user at the time of application.

Crop	Target Diseases	Use Rate fl. oz. product/A (lb. a.i./A)	Remarks
Safflower	Downy mildew (<i>Plasmopora halstedii</i> , <i>Plasmopora helianthi</i>) Alternaria leaf spot (<i>Alternaria</i> spp.)	6.0-15.5 (0.1-0.25)	<p>Integrated Pest (Disease) Management: Quadris should be integrated into an overall disease management strategy that includes varieties with disease tolerance and crop rotation to reduce plant debris in which inoculum overwinters.</p> <p>Resistance Management : Follow the resistance management guidelines in the Resistance Management Section. Do not apply more than two sequential applications of Quadris or other Group 11 fungicides before alternation with a fungicide that is not in Group 11.</p> <p>Application Directions: Apply 7.0 fl oz of Quadris at early bud followed by 14.0 fl. oz. at about 45 days before harvest. A third application of 7.0 fl. oz. may be made 30 days before harvest. Applications may be made by ground, air or chemigation. Use a minimum of 10 gallons of water per acre for ground applications.</p>
Specific Use Restrictions: Do not apply more than 27.5 fl.oz product/A (0.45 lb. a.i./A) per season of azoxystrobin-containing products. Do not apply within 30 days of harvest.			

Notes:

(1) It is extremely important to time the application of Quadris Flowable Fungicide appropriately as described above. Failure to do so may result in poor control. (2) Apply Quadris Flowable Fungicide in sufficient water volume to provide good coverage of the safflower head. (3) Syngenta does not recommend mixing this product with any other crop protection product.

Quadris® Flowable Fungicide is a registered trademark of Syngenta

©2006 Syngenta Crop Protection, Inc.

Registrant: Syngenta Crop Protection, Inc.
PO Box 18300
Greensboro, NC 27419

MT1098040AA0706