

---

**PRODUCT INFORMATION**

---



**Syngenta Crop Protection, Inc.**  
Greensboro, NC 27409  
www.syngenta-us.com

**RESTRICTED USE PESTICIDE**  
**Toxic to Fish Mammals, and Aquatic Organisms**

For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

**SUPPLEMENTAL LABEL FOR PROCLAIM® INSECTICIDE**

**INSECTICIDE**

**EPA Reg. No. 100-904**

**Supplemental Directions for Use on Fruiting Vegetables (except Cucurbits); *Brassica (Cole)* Leafy Vegetables; Leafy Vegetables (except *Brassica*); and Turnip Greens (tops, leaves).**

Active Ingredient:	
Emamectin benzoate (CAS No. 155569-91-8).....	5.0%
Other Ingredients:	95.0%
Total:	100.0%

Proclaim Insecticide is a water-dispersible granule containing 5% active ingredient.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**This labeling must be in the possession of the user at the time of pesticide application.**

SCP 904A-S1 0803

**DIRECTIONS FOR USE**

**Fruiting Vegetables (except Cucurbits): Tomato; Peppers: bell, chili, cooking, pimento, and sweet; Eggplant; Ground cherry; Pepino; Tomatillo.**

Pest	Rate Per Acre Per Application	Remarks	PHI
Beet armyworm Cabbage looper Fall armyworm Southern armyworm Tobacco budworm Tobacco hornworm Tomato hornworm Tomato fruitworm Tomato pinworm Yellowstriped armyworm	<b>Foliar application:</b>  2.4-4.8 oz. /A	Apply when larvae are first observed. Additional applications may be made to maintain control.  Use 2.4 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	7 days
Alfalfa looper Soybean looper <i>Liriomyza</i> leafminers <sup>1</sup>	<b>Foliar application:</b>  3.2-4.8 oz. /A	Apply when larvae are first observed. Additional applications may be made to maintain control.  Use 3.2 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	7 days

<sup>1</sup>Proclaim Insecticide provides suppression of *Liriomyza trifolii* and *Liriomyza sativae* populations. Suppression means either erratic control ranging from good to poor, or consistent control at a level below that which is generally considered acceptable for commercial control.

- **Do not** apply more than 4.8 oz. /A per application.
- **Do not** apply more than a total of 28.8 oz. /A per crop per season.
- Allow a minimum of 7 days between applications.
- Allow 7 days (PHI) between the last application and harvest.
- **Do not** allow livestock to graze in treated areas.

**Brassica Head and Stem Vegetables:** [Broccoli; Brussels sprouts; Cabbage; Cauliflower; Cavalo broccolo; Chinese broccoli (gai lon); Chinese (napa) cabbage ; Chinese mustard cabbage (gai choy); Kohlrabi] and

**Brassica Leafy Vegetables:** [Broccoli raab (rapini); Chinese (bok choy) Cabbage; Collards; Kale; Mizuna; Mustard greens; Mustard spinach; Rape greens], and Turnip greens (tops, leaves)<sup>2</sup>

Pest	Rate Per Acre Per Application	Remarks	PHI
Beet armyworm Cabbage webworm Corn earworm Cross-striped cabbageworm Diamondback moth Fall armyworm Imported cabbageworm	<b>Foliar application:</b>  2.4-4.8 oz. /A	Apply when larvae are first observed and repeat applications as necessary to maintain control.  Use 2.4 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	Brassica Head and Stem Vegetables: 7 days
Cabbage looper Soybean looper <i>Liriomyza</i> leafminers <sup>1</sup>	<b>Foliar application:</b>  3.2-4.8 oz. /A	Apply when larvae are first observed and repeat application as necessary to maintain control  Use 3.2 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	Brassica Leafy Vegetables and Turnip Greens: 14 days

<sup>1</sup> Proclaim Insecticide provides suppression of *Liriomyza trifolii* and *Liriomyza sativae* populations. Suppression means either erratic control ranging from good to poor, or consistent control at a level below that which is generally considered acceptable for commercial control.

<sup>2</sup>**Turnip Greens (tops, leaves) only:** For use on turnip varieties grown for leaves only. Do not use on turnip varieties grown for roots or dual-purpose varieties grown for roots and leaves.

- **Do not** apply more than 4.8 oz. /A per application.
- **Do not** apply more than a total of 28.8 oz. /A per crop per season.
- Allow a minimum of 7 days between applications.
- **Do not** allow livestock to graze in treated areas.

**Leafy Vegetables (except *Brassica*): Amaranth (leafy amaranth, Chinese spinach, tampala); Arugula (Roquette); Cardoon; Celery; Celtuce; Chervil; Chinese celery; Edible-leaved chrysanthemum; Garland chrysanthemum; Corn salad; Cress (garden and upland); Dandelion; Dock (sorrel); Endive (escarole); Florence fennel (finocchio); Head lettuce; Leaf lettuce; Orach; Parsley; Purslane (garden and winter); Radicchio (red chicory); Rhubarb; Spinach; New Zealand spinach; Vine spinach (Malabar spinach, Indian spinach); Swiss chard**

Pest	Rate Per Acre Per Application	Remarks	PHI
Beet armyworm Corn earworm Fall armyworm Tobacco budworm	<b>Foliar application:</b>  2.4-4.8 oz. /A	Apply when larvae are first observed and repeat application as necessary to maintain control.  Use 2.4 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	7 days
Cabbage looper Soybean looper <i>Liriomyza</i> leafminers <sup>1</sup>	<b>Foliar application:</b>  3.2-4.8 oz. /A	Apply when larvae are first observed and repeat application as necessary to maintain control.  Use 3.2 oz. /A for low to moderate infestations and 4.8 oz. /A for high infestations.	7 days

<sup>1</sup> Proclaim Insecticide provides suppression of *Liriomyza trifolii* and *Liriomyza sativae* populations. Suppression means either erratic control ranging from good to poor, or consistent control at a level below that which is generally considered acceptable for commercial control.

- Do not apply more than 4.8 oz. /A per application.
- Do not apply more than a total of 28.8 oz. /A per crop per season.
- Allow a minimum of 7 days between applications.
- Allow 7 days (PHI) between the last application and harvest.
- Do not allow livestock to graze in treated areas.

Proclaim® and the Syngenta logo are trademarks of a Syngenta Group Company  
 © 2003 Syngenta

Syngenta Crop Protection, Inc.  
 Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP 904A-S1 0803

Pro904A-S1 0803-Ig-8-8-03