
PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

SUPPLEMENTAL LABEL FOR QUILT™

FUNGICIDE

EPA Reg. No. 100-1178

Supplemental Directions For Applications on Pecans

Active Ingredient:	
Azoxystrobin	7.0%
Propiconazole	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Quilt as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR APPLICATIONS

Crop	Target Diseases	Use Rate fl. oz./product/A	Remarks
Pecans	Pecan scab (<i>Cladosporium caryigenum</i>) Anthracnose (<i>Glomerella cingulata</i>) Downy spot (<i>Mycosphaerella caryigena</i>) Liver spot (<i>Gnomonia caryae</i> pv <i>pecanae</i>) Vein spot (<i>Gnomonia nerviseda</i>) Zonate leaf spot (<i>Cristulariella moricola</i>) Powdery mildew (<i>Microsphaera penicillata</i>)	14.0–27.5	<p>Pecan Scab: Apply 14.0–27.5 fl. oz./A Quilt on a 14-day schedule during bud break and pre-pollination sprays. Apply 14.0–27.5 fl. oz./acre during nut formation and cover sprays. Use higher rates when disease pressure is heavier. Do not apply after shuck split.</p> <p>Other Foliar Diseases: Quilt may be applied for control of mid to late season foliar diseases at 14.0–20.5 fl. oz./A with other pecan products labeled for these diseases. Observe all directions, precautions, and limitations for the other products. Make no more than two sequential applications of a Group 11 fungicide before alternating to another product with a different mode of action than Group 11 fungicides. Ground applications apply in sufficient water to provide for full coverage. Aerial applications apply in a minimum of 20 gallons of water.</p>
<p>Specific Use Restrictions: For ground application, apply with enclosed cabs only. Do not apply more than 122 fl. oz./A of Quilt per season. Do not apply after shuck split or within 45 days of harvest, whichever is first. Do not graze livestock in treated areas or cut treated cover crops for feed. Do not apply more than 1.0 lb. ai propiconazole containing products/A/year. Do not apply more than 1.2 lb. ai azoxystrobin containing products/A/year.</p>			
<p>The active ingredient(s) in this product may have effects on federally listed threatened and endangered species or critical habitat in some counties. When using this product, you must follow the measures contained in the County Bulletin for the county in which you are applying the pesticide. To determine whether your County has a Bulletin consult http://www.epa.gov/espp/usa-map.htm. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.</p>			

Trademark of Syngenta

©2005 Syngenta

Syngenta Crop Protection, Inc.
 Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1178A-S3 0205

QUI 1178A-S3 0205 – bb – 2-17-05