
PRODUCT INFORMATION



Syngenta Crop Protection, Inc.
Greensboro, NC 27409
www.syngenta-us.com

SUPPLEMENTAL LABEL FOR QUILT FUNGICIDE

EPA Reg. No. 100-1178

For Disease Control in Cereals

Active Ingredient:	
Azoxystrobin	7.0%
Propiconazole	11.7%
Other Ingredients:	81.3%
Total:	100.0%

Contains 1.04 lbs. a.i. propiconazole and 0.62 lb. a.i. azoxystrobin per gallon.

CAUTION

See additional precautionary statements and directions for use inside booklet.

KEEP OUT OF REACH OF CHILDREN.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

Before using Quilt as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.

This supplemental label must be in possession of the user at the time of application.

DIRECTIONS FOR USE

Crop	Target Diseases	Use Rate fl. oz. product/A	Remarks
Cereals Barley Wheat Triticale	Early season suppression of: Powdery mildew <i>(Blumeria spp., Erysiphe spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Glume Blotch <i>(Stagonospora nordorum)</i> Tan spot <i>(Pyrenophora tritici-repentis)</i>	7-14	Apply Quilt in the spring for suppression of early season diseases. Applications may be made no closer than a 14-day interval. Quilt may be applied up to Feekes growth stage 9. Quilt may be applied by ground, air, or chemigation.
	Control of Leaf Diseases: Rust <i>(Puccinia spp.)</i> Powdery mildew <i>(Blumeria spp., Erysiphe spp.)</i> Leaf Blight <i>(Septoria tritici)</i> Glume Blotch <i>(Stagonospora nordorum)</i> Tan spot <i>(Pyrenophora tritici-repentis)</i> Helminthosporium leaf blight <i>(Drechslera tritici-repentis)</i> Spot Blotch <i>(Bipolaris sorokiniana)</i> Barley Scald <i>(Rhynchosporium secalis)</i> Barley stripe <i>(Pyrenophora graminea)</i> Net blotch <i>(Pyrenophora teres)</i> Kernel blight <i>(Alternaria spp.)</i>	10.5-14	Protecting the flag leaf is important for maximizing the potential yield. Highest yields are normally obtained when Quilt is applied when the flag leaf is 50% to fully emerged. If disease pressure is low, 10.5 fl. oz/acre may be applied. Applications may be made no closer than a 14-day interval. Quilt can be applied until the ligule of the flag leaf emerges (Feekes growth stage 9) on barley, rye, and triticale. For wheat only, Quilt can be applied until full head emergence (Feekes growth stage 10.5). Do not apply after this stage to avoid possible illegal residues. Quilt may be applied by ground, air or chemigation.
	Foot Rot/Eyespot <i>(Tapesia spp.)</i>	14	Apply full rate of Quilt plus half the rate recommended of other EPA registered fungicides such as Topsin® M. Apply at tillering but before elongation has occurred.
Specific Use Restrictions: Do not apply more than 2 applications/A/year. Do not apply within 45 days of harvest for grain and straw. Do not harvest wheat for forage. Do not graze or feed livestock treated forage or cut green crop for hay or silage. Do not apply more than 20.5 fl. oz./A/season of Quilt. Do not apply more than 0.167 lb ai propiconazole containing products/A/year. Do not apply more than 0.4 lb ai azoxystrobin containing products/A/ year. Under certain environmental conditions, tank mixes of Quilt plus herbicides and/or fertilizers may cause crop injury in barley, triticale and wheat.			

Trademark of Syngenta

©2005 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, North Carolina 27409
www.syngenta-us.com

SCP 1178A-S5 0605

Qui 1178A-S5 605 nsp 06/28/05