

---

**SUPPLEMENTAL LABEL**

---



**Syngenta Crop Protection, Inc.**  
P. O. Box 18300  
Greensboro, North Carolina 27419-8300

**Scholar® SC**

**Fungicide**

Active Ingredient:	
Fludioxonil:*	20.4%
Other Ingredients:	79.6%
Total:	100.0%

\*CAS No. 131341-86-1

Scholar SC is a flowable suspension concentrate

Scholar SC contains 1.92 lbs. ai per gallon

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

EPA Reg. No. 100-1242

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**Before using Scholar SC as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions and precautions on all EPA-registered products are to be followed. This label and the Federally registered label must be in the possession of the user at the time of pesticide application.

---

### CROP USE DIRECTIONS

---

#### Sweet Potato

Use Scholar SC as a post-harvest dip and low volume application for the control of post-harvest rots caused by *Rhizopus stolonifer*.

Application Method	Disease	Rate (fl. oz.)	Remarks
In-Line Dip/Drench	Rhizopus rot	16-32 fl. oz./100 gals.	<ul style="list-style-type: none"> <li>Mix 16-32 fl. oz. of Scholar SC in 100 gals. of water, wax/emulsion, or aqueous dilution of wax/oil emulsion.</li> <li>Dip for approximately 30 seconds and allow fruit to drain.</li> <li>Add 8 fl. oz. of Scholar SC to 100 gals. of treating suspension after 500 bushels are treated.</li> <li>After each 1000 bushels treated, drain and flush the tank. Refill with fresh dip suspension.</li> </ul>
In-line Aqueous or Fruit Coating Spray application	Rhizopus rot	16 fl. oz./200,000 lbs. of sweet potatoes	<ul style="list-style-type: none"> <li>Ensure proper coverage of the fruit.</li> <li>Mix 16 fl. oz. of Scholar SC in an appropriate water, wax/oil emulsion, or aqueous dilution of a wax/oil emulsion for the crop being treated.</li> <li>Use T-Jet, CDA, or similar application system.</li> </ul>
<b>Do not make more than one post-harvest application to the sweet potatoes.</b>			
<ul style="list-style-type: none"> <li>Ensure the Scholar SC solution remains in suspension by using agitation.</li> <li>Scholar SC is stable in chlorine (100 ppm solution) and at temperatures of 60°C (or 140°F) that can be used to disinfest high-volume, recycling tanks.</li> </ul>			

**NOTE:** Ensure the Scholar SC solution remains in suspension by using agitation. Scholar SC may be degraded by exposure to direct sunlight. Treated sweet potatoes should not be stored in direct sunlight.

---

Manufactured for:  
Syngenta Crop Protection, Inc.  
P.O. Box 18300  
Greensboro, North Carolina 27419-8300

SCP 1242A-S2 0209

Scholar SC 1242A-S2 0209 - bb - 7-9-09