

---

**PRODUCT INFORMATION**

---



**Syngenta Crop Protection, Inc.**  
Greensboro, NC 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

**SUPPLEMENTAL LABEL FOR SWITCH 62.5WG**

**FUNGICIDE**

**EPA Reg. No. 100-953**

For control of diseases on berries, pistachio and watercress

**Active Ingredients:**

Cyprodinil (CAS No. 121552-61-2) ..... 37.5%

Fludioxonil (CAS No. 131341-86-1) ..... 25.0%

Other Ingredients: ..... 37.5%

**Total:** ..... 100.0%

Switch 62.5WG is a water-dispersible granule.

**KEEP OUT OF REACH OF CHILDREN.**

**CAUTION**

**All applicable directions, restrictions and precautions on the EPA-registered label are to be followed.**

**This labeling must be in the possession of the user at the time of pesticide application.**

SCP 953A-S1 0303

## GENERAL INFORMATION

Do not apply this product with fixed wing or rotary aircraft.

### Chemigation

Do not apply this product through any type of irrigation system.

## USE DIRECTIONS FOR SPECIFIC CROPS

**Rotational Crop Restrictions:** Do not plant any other crop for a period of 12 months unless Switch 62.5WG is registered for that use.

Crop	Disease	Rate oz./Acre	Application Timing	Notes
<b>Berry Crop Grouping:</b>  <b>Bushberries</b> (blueberries, currants, gooseberries, elderberry, huckleberry, as well as other cultivars and hybrids of these).  <b>Caneberries</b> (blackberries, raspberries, and/or cultivars and/or hybrids of these).  <b>Juneberry</b> <b>Lingonberry</b> <b>Salal</b>	Mummy Berry ( <i>Monilinia vaccinii-corymbosi</i> )	11-14	Make the first application during early bloom. A second application should be made 7 to 10 days later. Additional applications can be made at 7-10 day intervals if conditions remain favorable for disease development.	Pre-harvest interval is 0 days.  Do not apply more than 56 ounces of product per acre per year.
	Anthracnose ( <i>Colletotrichum spp.</i> )			
	Alternaria Fruit Rot ( <i>Alternaria tenuissima</i> )			
	Phomopsis ( <i>Phomopsis vaccinii</i> )		Make no more than two (2) sequential applications before using another registered fungicides.	
	Botrytis Fruit Rot ( <i>Botrytis cinerea</i> )		Switch 62.5WG may be applied in an alternating or blocking program.	

Crop	Disease	Rate oz./Acre	Application Timing	Notes
<b>Pistachio</b>	<p>Botrytis (<i>Botrytis</i>)</p> <p>Alternaria (<i>Alternaria</i>)</p>	11-14	<p>Make the first application during early bloom. A second application should be made 14 days later. Additional applications can be made at 14 day intervals if conditions remain favorable for disease development.</p> <p>Make no more than two (2) sequential applications before using another registered fungicide. Switch 62.5WG may be applied in an alternating or blocking program.</p>	<p>Do not apply Switch 62.5WG within 7 days of harvest. (7 day PHI)</p> <p>Do not apply more than 56 ounces of product per acre per year.</p>
<b>Strawberries</b>	Gray Mold ( <i>Botrytis cinerea</i> )	11-14	<p>Begin application at or before bloom and continue on a 7 to 10-day interval.</p> <p>Make no more than two (2) sequential applications before using another registered fungicide. Switch 62.5WG may be applied in an alternating or blocking program .</p>	<p>Preharvest interval is 0-days.</p> <p>Do not exceed 56 ounces of product per year per acre.</p>
<b>Watercress</b>	<p>Cercospora leafspot (<i>Cercospora spp.</i>)</p> <p>Sclerotinia white mold (<i>Sclerotinia spp.</i>)</p> <p>Rhizoctonia rot (<i>Rhizoctonia solani</i>)</p>	11-14	<p>Begin applications when disease first appears and continue on a 7-10 day interval.</p> <p>Make no more than two (2) sequential applications before using another registered fungicides. Switch 62.5WG may be applied in an alternating or blocking program.</p>	<p>Pre-harvest interval is 0 days.</p> <p>Do not apply more than 56 ounces of product per acre per year.</p>

Switch™ and the Syngenta logo are trademarks of a Syngenta Group  
Company  
U.S. Patent No.5,153,200

© 2003 Syngenta

Syngenta Crop Protection, Inc.  
Greensboro, North Carolina 27409  
[www.syngenta-us.com](http://www.syngenta-us.com)

SCP953A-S1 0303

SWI 62.5WG 953A-S1 0303 supp label - nsp – 3-17-03, revised 4-28-03 aa