
SUPPLEMENTAL LABELING

Syngenta Crop Protection, Inc.
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1169B-S5 0210

GROUP	9	HERBICIDE
-------	----------	-----------

Traxion®

Herbicide

Nonselective Foliar Systemic Herbicide for Weed Control

Active Ingredient:

*Glyphosate: N-(phosphonomethyl) glycine 36.5%

Other Ingredients: 63.5%

Total: 100.0%

*Contains 500 grams per liter or 4.17 pounds per U.S. gallon of glyphosate acid.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1169

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Traxion as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

SUGAR BEET, GLYPOSATE TOLERANT

For use on the following glyphosate tolerant sugarbeet only: all Roundup Ready® sugarbeet including Roundup Ready sugarbeet, and all Genuity™ brand sugarbeet which contains Roundup Ready.

Method of Application: Before, during, or after planting; and postemergence in sugarbeet varieties which have been genetically modified to be tolerant to glyphosate based herbicides.

Follow directions listed in the **APPLICATION DIRECTIONS, SPRAY ADDITIVES**, and **APPLICATION EQUIPMENT AND TECHNIQUES** sections. Refer to **WEEDS CONTROLLED** section for application rates and timing.

Use Precautions and Restrictions for Glyphosate Tolerant Sugarbeet

- Maximum amount of Traxion which may be applied from all applications is 5.7 qt/A.
- Maximum preplant/preemergence rate is 3.6 qt/A.
- Maximum amount of Traxion applied over-the-top of the crop from emergence to 8-leaf state is 60 fl oz/A.
- From emergence to 8-leaf stage apply no more than 35 fl oz/A for any single application.
- Maximum amount of Traxion applied over-the-top of the crop from 8-leaf state to canopy closure is 48 fl oz/A.
- From 8-leaf state to canopy closure apply no more than 24 fl oz/A for any single application.
- Make no more than 4 sequential postemergence applications of Traxion, which must be at least 10 days apart.
- Allow a minimum of 30 days between postemergence application and harvest.
- Applications of Traxion to sugarbeet varieties which are not glyphosate tolerant will result in severe crop injury and reduced yields.

Traxion® and the Syngenta logo are trademarks of a Syngenta Group Company. Roundup Ready® and Genuity™ are trademarks of Monsanto Technology LLC.

©2010 Syngenta