



ETIRA

Section 24(c) Special Local Need Label

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF
SOUTH CAROLINA**

Reflex® Herbicide

EPA Reg. No. 100-993

EPA SLN No. SC-110002

Early Preplant Application on Medium and Fine-Textured Soils for Weed Control in Cotton

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY
RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered Reflex label and this accompanying supplemental label. This labeling must be in the possession of the user at the time of pesticide application.

Cotton

Apply Reflex at 1 pt/A to medium or fine-textured soils (i.e. soil types heavier than coarse textured soils) as a preplant surface application up to 21 days prior to planting cotton, and after the last tillage operation is completed. A minimum of 0.5 inch of rainfall or overhead irrigation must occur before planting cotton. Reflex will provide preemergence control of Palmer amaranth (including glyphosate-resistant Palmer amaranth) pigweed species, and control or partial control of other broadleaf weeds, sedges and grasses listed on the federal label.

To broaden the weed control spectrum, Reflex may be tank mixed with other preemergence herbicides such as Caparol®, Cotoran®, Direx®, Karmex®, Solicam®, or Staple®. For control of emerged weeds, Reflex may be tank mixed with a burndown herbicide such as dicamba, Gramoxone Inteon™ or glyphosate brands (such as Touchdown®, Roundup®) labeled in cotton. Refer to the tank-mix partner label for use directions, restrictions and limitations. The most restrictive product labeling applies.

Cotton plants are tolerant to preplant applications of Reflex when applied at the recommended rate and application use directions. Some crinkling or spotting of cotton foliage or stunting may occur, but cotton plants normally outgrow these effects and develop normally.

General Use Directions, Restrictions and Precautions

- After Reflex application, a minimum of 0.5 inch of rainfall or overhead irrigation must occur before planting cotton on medium or fine-textured soils. Failure to follow these use directions will result in severe crop injury.
- Do not disturb or re-work the seedbed following a Reflex application.
- Cotton must be planted at least 0.75 inch in depth.
- Avoid overlapping spray swaths.
- The use of an in-furrow or seed applied fungicide will generally assist with seedling establishment and development.

Caparol®, Gramoxone Inteon™, Reflex®, Solicam®, Touchdown® trademarks of a Syngenta Group Company.

Direx®, Karmex®, Staple® trademarks of E. I. du Pont de Nemours & Co., Inc.

Roundup® trademark of Monsanto Company.

Cotoran® trademark of Agan Chemical Manufacturing, Ltd.

®Reflex trademark of a Syngenta Group Company

© 2011 Syngenta Crop Protection, Inc.

24(c) registrant:
Syngenta Crop Protection
Greensboro, NC 27419-8300

Label Code: SC0993021AA0411