
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 620A-S2 0110

Caparol® 4L

Herbicide

Active Ingredient:	
Prometryn: 2,4-bis(isopropylamino)- 6-(methylthio)-s-triazine	44.4%
Other Ingredients:	55.6%
Total:	100.0%

Caparol 4L contains 4 lbs. active ingredient per gallon.

Shake well before using.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-620

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Caparol 4L as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



DIRECTIONS FOR USE

CROP USE DIRECTIONS

Carrot

Caparol 4L can be applied preemergence and or postemergence over the top to carrot. Read and follow all directions for use in carrot.

Use Precautions for Pre and Postemergence Applications in Carrot

Apply up to three applications of Caparol 4L at the rate of 2 - 4 pt/A per application.

Do not exceed one preemergence at up to 4 pt/A and two postemergence applications each at up to 2 pt/A or one postemergence application at up to 4 pt/A per crop cycle.

Do not exceed 8 pt/A of Caparol 4L per crop cycle.

Make postemergence applications through the 6 leaf stage of carrot development.

Do not apply within 30 days of harvest.

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

When applying to emerged weeds add 2 qt of a nonionic surfactant (NIS) or wetting agent (approved for intended use) to 100 gal of spray mixture (0.5%) v/v or 1 gal of a non-phytotoxic crop oil concentrate (COC) containing 15-20% approved emulsifier to 100 gal of spray mixture (1% v/v).

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on carrots: cabbage, carrots, leafy petioles crop subgroup 4B, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Celeriac

Do not exceed 4 pt/A of Caparol 4L per crop cycle.

Transplants

Make a single broadcast application at 1.6-4 pt/A in a minimum of 20 gal of water per acre after the crop has 6-8 leaves. Application may be made over the crop. Within the rate ranges given, use the lower rate on relatively coarse-texture soils and soils low in organic matter; use the higher rate on relatively fine-textured soils and soils high in organic matter. Apply before weeds are 2 inches tall. Do not apply to the crop within 60 days of harvest.

Cilantro

Caparol 4L can be applied post planting, preemergence to cilantro. Read and follow all directions for use in cilantro.

Use Precautions for Preemergence Applications in Cilantro

Make one application of Caparol 4L at the rate of 2 – 3.2 pt/A.

Use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and soils high in organic matter.

Do not exceed 3.2 pt/A of Caparol 4L per crop cycle.

Do not apply within 30 days of harvest.

Make uniform applications of the herbicide in a minimum of 20 gallons of water per acre.

Do not use on sand or loamy sand soil.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 3.4 pt Caparol 4L per acre on cilantro: cabbage, carrots, leafy petioles crop subgroup 4B, cilantro, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Okra

Apply Caparol 4L preemergence and/or post-directed to okra. Read and follow all directions for use in okra.

Use Precautions for Pre and Post-Directed Applications in Okra

Make a single preemergence broadcast application of Caparol 4L after planting before crop emergence at the rate of 3.0 pt/A.

For two applications make the first preemergence after planting, before crop emergence at the rate of 1.5 pt/A and the second post-directed when okra plants are at 7-9 leaf stage at the rate of 1.5 pt/A. Apply post-directed before weeds are 2 inches tall.

Do not apply within 14 days of harvest.

Do not exceed one preemergence and one post-directed application per crop cycle.

Do not exceed 3 pt/A of Caparol 4L per crop cycle.

Make uniform applications of the herbicide in 20-40 gallons of water per acre.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on okra: cabbage, carrots, leafy petioles crop subgroup 4B, corn, cotton, dill, fennel, okra and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Parsley

In California only, make a single preemergence broadcast application of Caparol 4L after planting before crop emergence at the rate of 1.0-4.0 pt/A. If a rate higher than 1.0 pt/A is applied, then only one application is allowed.

In all states (including California), make a single preemergence broadcast application of Caparol 4L up to 14 days after planting at the rate of 1.0 pt/A. For extended weed control, a second application can be made at 1.0 pt/A up to 30 days prior to harvest. A third application at 1.0 pt/A can be made to the regrowth up to 30 days prior to the second (cutting) harvest.

Do not make more than 3 application per year at a maximum of 1 pt/A per application except in California only where a single application up to 4 pt/A is allowed. Do not apply to the crop within 30 days of harvest.

Use the lower rates on coarse-textured soils and soils low in organic matter; use the higher rates on fine-textured soils and soils high in organic matter. Make a uniform application of the herbicide in 20 gallons of water per acre.

Precautions: To avoid injury to parsley, (1) Do not use on sand or loamy sand. (2) Do not apply if parsley is under water stress.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on okra: cabbage, carrots, leafy petioles crop subgroup 4B, corn, cotton, dill, fennel, okra, and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Rhubarb

Make a single broadcast application at the appropriate rate to established rhubarb when plants are dormant, before leaves have emerged from the crown. Apply 2.0-3.2 pt/A on coarse-textured soils and 3.2-4.0 pt/A on fine-textured soils. Apply in a minimum of 20 gallons of water per acre.

Within the rate ranges given, use the lower rate on relatively coarse-textured soils and soils low in organic matter; use the higher rate on relatively fine-textured soils and soils high in organic matter.

Do not apply to rhubarb within 40 days of harvest.

Rotational Crops

The following crops may be seeded 5 months after applying no more than 4 pt Caparol 4L per acre on rhubarb: cabbage, carrots, leafy petioles crop subgroup 4B, corn, cotton, dill, fennel, okra, and peas. Onions and red beets may not be planted within 8 months of applying Caparol 4L. All other crops may be planted 12 months after applying Caparol 4L.

Caparol® and the Syngenta logo are trademarks of a Syngenta Group Company

©2011 Syngenta