
SUPPLEMENTAL LABELING

Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, North Carolina 27419-8300
SCP 1070A-S2 0111

FUSILADE® DX

GROUP	1	HERBICIDE
-------	---	-----------

HERBICIDE

Postemergence Herbicide for Control of Perennial and Annual Grass Weeds

Active Ingredient:

Fluazifop-P-butyl

Butyl (*R*)-2-[4-[[5-(trifluoromethyl)-2-pyridinyl]oxy]phenoxy]propanoate* 24.5%

Other Ingredients: 75.5%

Total: 100.0%

*Fusilade DX Herbicide contains 2 pounds (+) isomer (fluazifop-P-butyl) per gallon.

Contains petroleum distillates.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

EPA Reg. No. 100-1070

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using Fusilade DX as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

syngenta

DIRECTIONS FOR USE

CROP SPECIFIC RESTRICTIONS AND LIMITATIONS

BANANAS AND PLANTAINS

- Do not apply a total of more than 72 fl. oz. of Fusilade DX Herbicide per acre per season to bananas or plantains.
- Do not apply more than 24 fl. oz. of Fusilade DX Herbicide per acre per application.
- Maintain a minimum of 30 days between applications.
- Bananas or plantains may be harvested 0 days after the last application.

CITRUS FRUITS - Calamondin, Citrus citron, Citrus hybrids, Grapefruit, Kumquat, Lemon, Lime, Mandarin (Tangerine), Orange (all), Pummelo, and Satsuma mandarin

- Do not apply a total of more than 72 fl. oz. of Fusilade DX Herbicide per acre per season to citrus fruit crops.
- Do not apply more than 24 fl. oz. of Fusilade DX Herbicide per acre per application.
- Maintain a minimum of 21 days between applications.
- Do not harvest citrus fruit within 14 days of last application.

GRAPES

- Do not apply a total of more than 72 fl. oz. of Fusilade DX Herbicide per acre per season to grapes.
- Do not apply more than 24 fl. oz. of Fusilade DX Herbicide per acre per application.
- Maintain a minimum of 14 days between applications.
- Do not harvest grapes within 50 days of last application.

SUGAR BEETS

- Do not apply a total of more than 48 fl. oz. of Fusilade DX Herbicide per acre per season to sugar beets.
- Do not apply more than 24 fl. oz. of Fusilade DX Herbicide per acre per application.
- Maintain a minimum of 14 days between applications.
- Do not harvest sugar beets within 90 days of last application.

SPECIAL INSTRUCTIONS FOR USE IN BANANAS AND PLANTAINS

Apply Fusilade DX Herbicide as a directed postemergence application in the interspaces and around the base of banana and plantain plants while avoiding contact of the spray solution or mist with the trunk and foliage. Applications can be made as a broadcast, strip band, or spot spray application at rates and growth stages listed in

Tables 2-5. Apply with sufficient spray volume and pressure to ensure complete coverage of target grasses.

SPECIAL INSTRUCTIONS FOR USE IN APRICOTS, CHERRIES, CITRUS FRUITS [CALAMONDIN, CITRUS CITRON, CITRUS HYBRIDS, GRAPEFRUIT, KUMQUAT, LEMON, LIME, MANDARIN (TANGERINE), ORANGE (ALL), PUMMELO, AND SATSUMA MANDARIN], GRAPES, MACADAMIA NUTS, NECTARINES, PEACHES, PECANS, PLUMS AND PRUNES

Apply Fusilade DX Herbicide as a directed postemergence application in the interspaces and around the base of the trees while avoiding contact of spray with foliage of trees. Apply with sufficient spray volume and pressure to ensure complete coverage of target grasses. Applications can be made as a broadcast, strip band, or spot spray application at rates and growth stages listed in Tables 2-5.

Fusilade® DX and the Syngenta logo are trademarks of a Syngenta Group Company.

Manufactured for:
Syngenta Crop Protection, LLC
P.O. Box 18300
Greensboro, North Carolina 27419-8300

SCP 1070A-S2 0111

Fusilade DX 1070A-S2 0111 - bb - 4-27-11