

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Entrust[®]

EPA Reg. No. 62719-282

**For Control of Armyworms in Legume Forages and Legume Forage Grown for Seed
(including but not limited to Alfalfa, Clover, Lupin, and Vetch)**

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Entrust[®] Naturalyte[®] insect control before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Entrust according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Entrust.

Directions for Use

Refer to product label for General Use Precautions, Mixing and Application Instructions.

Pests and Application Rates:

Pests	Rate (oz/acre)	Active Ingredient (lb/acre)
beet armyworm fall armyworm southern armyworm true armyworm	0.63 – 1.25	0.031 – 0.062

Specific Use Directions:

Application Timing: Scout at least weekly and consider the impact of both pests and beneficials. Treat when economic thresholds are exceeded, targeting eggs at hatch or small larvae. Heavy infestations may require repeat applications, but follow resistance management guidelines on the product label. Consult your Dow AgroSciences representative, extension service specialist, certified crop advisor or your state agricultural experiment station for any additional recommendations applicable to your area.

Application Rate: Use a higher rate in the rate range for heavy infestations or advanced growth stages of target pests.

Resistance Management: For resistance management, do not apply Entrust more than 3 times in any 21-day period. Rotate to a different class of insect control products or use no treatment for the next 21 days. Do not apply more than 6 treatments per season. If Entrust is applied 3 times in succession, do not apply again for at least 21 days.

Restrictions:

- **Preharvest Interval:** Do not harvest hay or fodder for 3 days after treatment. There is no preharvest interval for forage.
- Do not allow cattle to graze from treated area until spray has dried.
- Do not apply more than 3.75 oz of Entrust (0.186 lb ai spinosad) per acre per season.

Note: This product is toxic to bees exposed to treatment for 3 hours following treatment and toxic to aquatic invertebrates. Refer to the Environmental Hazards section of the product label attached to the product container for required protective measures.

[®]Trademark of Dow AgroSciences LLC

R184-004

EPA-accepted 02/24/06

Initial printing.