



## Supplemental Label

### For use on alfalfa, including alfalfa grown for seed

EPA Reg. No. 7969-199

#### Active Ingredients:

<b>pyraclostrobin:</b> (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester) .....	12.8%
<b>boscalid:</b> 3-pyridinecarboxamide, 2-chloro-N-(4'-chloro(1,1'-biphenyl)-2-yl)- .....	25.2%
<b>Other Ingredients:</b> .....	62.0%
<b>Total:</b> .....	100.0%

0.128 oz (0.008 lb) of pyraclostrobin in 1 oz of **Pristine®** fungicide  
0.252 oz (0.0158 lb) of boscalid in 1 oz of **Pristine**

Refer to the **Pristine** main label, EPA Reg. No. 7969-199, for complete **Directions For Use** and all applicable restrictions and precautions. This supplemental label and the full **Pristine** container label must be in the possession of the user at the time of pesticide application.

#### Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

#### General Information

**Pristine** provides optimum disease control when applied in a regularly scheduled protective fungicide program and used in a spray program that rotates fungicides with different modes of action. Refer to the **Pristine** main container label for general resistance management information and to the crop-specific use directions and restrictions found in this label.

Livestock grazing and feeding is permitted for crops listed in **Table 1. Pristine® fungicide Crop-specific Use Directions** in this label.

#### Application Information

Apply **Pristine** according to the rate, timing, resistance management, and adjuvant use recommendations in **Table 1. Pristine® fungicide Crop-specific Use Directions** in this label.

**Pristine** may be applied by ground sprayer, by aerial equipment, or through sprinkler irrigation systems. Refer to the **Pristine** main container label for specific instructions on these methods.

#### Restrictions and Limitations

**No aerial application in New York State except as permitted under FIFRA Section 24(c), Special Local Need Registration.**

**Pristine** is not for use in greenhouse or transplant production systems.

Follow the restrictions and limitations outlined in **Table 2. Pristine Restrictions and Limitations** in this label for:

- Minimum preharvest interval (PHI)
- Maximum product rate per acre
- Maximum number of applications
- Maximum product rate per season
- Livestock grazing or feeding

**Table 1. Pristine® fungicide Crop-specific Use Directions**

Crop	Target Disease	Product Use Rate per Application	Maximum Number of Applications per Season	Maximum Product Rate per Season	Minimum Time from Application to Harvest (PHI)
<b>Alfalfa (including alfalfa grown for seed)</b>	Anthracnose ( <i>Colletotrichum trifolii</i> )	12 to 18 ozs per acre	2 per cutting and 3 total per growing season	54 ozs per acre	14 days
	Common leaf spot ( <i>Pseudopezizza medicaginis</i> )				
	Downy mildew ( <i>Peronospora trifoliorum</i> )				
	Leaf spot ( <i>Leptosphaerulina briosiani</i> )				
	Powdery mildew ( <i>Erysiphe pisi</i> )				
	Rhizoctonia blight/black patch ( <i>Rhizoctonia</i> spp.)				
	Rust ( <i>Uromyces</i> spp.)				
	Spring black stem and leaf spot ( <i>Phoma medicaginis</i> )				
	Stagnospora leaf spot ( <i>Stagnospora meliloti</i> )				
	Stemphyllium leaf spot ( <i>Stemphyllium</i> spp.)				
	Summer black stem and leaf spot ( <i>Cercospora medicaginis</i> )				
	Yellow leaf blotch ( <i>Leptotrichila medicaginis</i> )				
	White mold/Sclerotinia crown and stem rot ( <i>Sclerotinia sclerotiorum</i> , <i>S. trifoliorum</i> )	14 to 18 ozs per acre			
	<b>Suppression only</b> Southern blight ( <i>Sclerotium rolfsii</i> )				

**Table 1. Pristine® fungicide Crop-specific Use Directions** *(continued)*

**Application Directions:** Begin **Pristine** applications when conditions favorable for disease are expected, but prior to onset of disease development. For stand establishment of fall seeded alfalfa, begin applications in fall through early winter prior to first snowfall or extended cool, wet conditions. For seed pod protection, begin applications at 10% to 30% bloom.

Using higher rates may improve disease control performance as the crop canopy volumes and densities increase. Disease control can also be improved when application equipment and spray volumes are adjusted to achieve thorough canopy penetration and coverage.

Repeat application on a 14-day to 21-day interval if conditions are conducive for disease development. **DO NOT** make more than two (2) applications per cutting or more than three (3) applications per season. Use the higher rate and shorter interval when disease pressure is high.

Under some conditions, the use of additives or adjuvants may improve the performance of **Pristine**.

**Resistance Management:** To limit the potential for development of resistance, **DO NOT** make more than two (2) applications of **Pristine per cutting or three (3) applications** per season.

**Table 2. Pristine Restrictions and Limitations**

Crop	Minimum Time from Application to Harvest (PHI) (days)	Maximum Product Rate per Acre per Application (ozs)	Maximum Number of Applications per Season	Maximum Product Rate per Acre per Season (ozs)	Livestock Grazing or Feeding Permitted
Alfalfa (including alfalfa grown for seed)	14	18	3	54	Yes

### Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.**

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.**

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. 1108

***Pristine** is a registered trademark of BASF.*

© 2008 BASF Corporation  
All rights reserved.

007969-00199.20081119.NVA 2008-04-156-0338

BASF Corporation  
26 Davis Drive  
Research Triangle Park, NC 27709

  
The Chemical Company