

Restricted Use Pesticide. Due to toxicity to aquatic invertebrate animals. For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS



Micromite® 80WGS

Insect Growth Regulator

EPA REG. NO. 400-487 EPA SLN NO. TX-110007

Active Ingredient: (% by weight)

Diflubenzuron

N-[[[4-Chlorophenyl)amino]carbonyl]-2,6-difluorobenzamide.....80%

Inert Ingredients:.....20%

Total:.....100%

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions on this supplemental label and the main EPA registered label. This labeling and the federal label for this product must be in the possession of the user at the time of pesticide application.

For application as Low Volume Ground Application:

Crop	Pest	Product Rate (oz./A)	Directions for Use
Grapefruit Sweet Orange Tangerine Pummelos/Pomelos and their hybrids	Asian Citrus Psyllid (nymphs & eggs)	6.25 oz. (2 pouches)	Apply in 2.0 gallons of finished spray solution per acre by ground using air-blast or air-assisted spray equipment. Spray nozzles that produce a droplet size with a volume median diameter of 90 microns or larger are required. Apply to immature citrus leaf flushes when Asian citrus psyllids are first observed in trees. Thorough coverage of foliage must be obtained for best control. Micromite 80WGS will not kill adult stages of Asian citrus psyllid.

Restrictions: Spray last three rows windward of surface water using nozzles on one side only, with spray directed away from surface water. Avoid spray going over tops of trees by adjusting or turning off top nozzles. Shut off nozzles on the side away from the grove when spraying the outside row. Shut off nozzles when turning at ends of rows and passing tree gaps in rows.

Do not make more than one low volume application per crop season. Rotate with other chemistries for resistance management.

Do not apply within 21 days of harvest.

Do not harvest cover crops for animal feed or graze livestock in treated groves.

Refer to label for additional restrictions and use directions