

Product Bulletin



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Delegate[®] WG

EPA Reg. No. 62719-541

Special 2(ee) Recommendation[†]
(For Distribution and Use Only in the State of Texas)

Low Volume Application in Citrus Fruits¹

ATTENTION

[†]This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Read the label affixed to the container for Delegate[®] WG insecticide before applying. Carefully follow all precautionary statements and applicable use directions except as specified below.
- Use of Delegate WG according to this Section 2(ee) recommendation is subject to all use precautions and limitations imposed by the label affixed to the container for Delegate WG.

Directions for Use

¹Citrus fruits (crop group 10) including calamondin, chironja, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, satsuma mandarin, sour orange, sweet orange, tangelo, tangor

Refer to product label for Delegate WG for General Use Precautions, Mixing Directions and Application Directions.

Apply Delegate WG by ground application in a minimum finished spray carrier volume of 2 gallons per acre. Spray applications must be made with spray nozzles that create a droplet size with a volume median diameter (VMD) of 90 microns or larger. An adjuvant may be used in the spray mixture at the specified rate of the manufacturer. Evaluate the effect of specific adjuvants or tank mix partners on the performance of Delegate WG in low volume applications in a small-scale application before making large-scale applications.

To achieve good control, uniform coverage and proper equipment calibration are required. Apply only with air-assisted equipment. Low volume applications may not provide the same level of efficacy that can be obtained with an airblast sprayer at higher application volumes.

Reduce the potential for drift when making low volume applications. Factors that may impact drift include wind speed and temperature inversion. Only apply this product if the wind direction favors on-target deposition. Do not apply when wind velocity exceeds 15 mph. See the Application Directions section for additional information on reducing spray drift.

[®]Trademark of Dow AgroSciences LLC

R359-019

Issued: 09/27/11

Initial printing