#### SUPPLEMENTAL LABELING

# RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

## Syngenta Crop Protection, LLC

P. O. Box 18300 Greensboro, North Carolina 27419-8300 SCP 1217B-S19 0711

#### Gramoxone® SL

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide Vegetables (Vine Desiccation) – For Use Only in the States of Delaware and Georgia

#### **Active Ingredient:**

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride)	30.1%
Other Ingredients:	69.9%
Total:	100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon. Contains alerting agent (odor), emetic, dye.

EPA Reg. No. 100-1217

### KEEP OUT OF REACH OF CHILDREN.



# DANGER / POISON PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.
- IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.
- DO NOT USE OR STORE IN OR AROUND THE HOME.
- DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.
- THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.



# **DIRECTIONS FOR USE**

Сгор	Maximum Number of Applications	Use Pattern	Gramoxone Inteon Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Broomstions Bostrictions and Comments
	Applications				interval	Precautions, Restrictions and Comments
VEGETABLES	2	Broadcast Spray Post-Harvest	2.25-3 pts.	Ground: 10 gals.		<ul> <li>For post-harvest desiccation of vegetable vines.</li> </ul>
(vine desiccation)						<ul> <li>Apply as a broadcast spray over-the-top of plants following harvest.</li> </ul>
For Use Only in the States of Delaware and Georgia						<ul> <li>Add a nonionic surfactant at 1 qt. per 100 gals. of spray mix.</li> <li>Do not graze livestock in treated area.</li> <li>Do not apply this product through any type of irrigation system.</li> </ul>

Gramoxone® SL and the Syngenta logo trademark of a Syngenta Group Company

©2011 Syngenta

GRAMOXONE SL 1217B-S19 0711 - tb - 08/24/11