

---

## SUPPLEMENTAL LABELING

---

### RESTRICTED USE PESTICIDE DUE TO ACUTE TOXICITY

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

#### **Syngenta Crop Protection, LLC**

P. O. Box 18300

Greensboro, North Carolina 27419-8300

**SCP 1217B-S2 0711**

#### **Gramoxone® SL**

A Weed, Grass, and Harvest Aid Desiccant/Defoliant Herbicide

Peanuts – For Use Only in the States of Florida and Georgia

Active Ingredient:

Paraquat dichloride (1,1'-dimethyl-4,4'-bipyridinium dichloride) ..... 30.1%

Other Ingredients: ..... 69.9%

Total: ..... 100.0%

Contains 2.0 pounds paraquat cation per gallon as 2.762 pounds salt per gallon.

Contains alerting agent (odor), emetic, dye.

**EPA Reg. No. 1217**

**KEEP OUT OF REACH OF CHILDREN.**



**DANGER / POISON**  
**PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

- **NEVER PUT INTO FOOD, DRINK OR OTHER CONTAINERS.**
- **IF SWALLOWED, TAKE IMMEDIATE ACTION AS PRESCRIBED IN FIRST AID STATEMENT. SYMPTOMS ARE PROLONGED AND PAINFUL.**
- **DO NOT USE OR STORE IN OR AROUND THE HOME.**
- **DO NOT REMOVE CONTENTS EXCEPT FOR IMMEDIATE USE.**
- **THE ODOR OF THIS PRODUCT IS FROM THE ALERTING AGENT WHICH HAS BEEN ADDED, NOT FROM PARAQUAT.**

All applicable directions, restrictions and precautions on the EPA-registered label are to be followed. Before using GRAMOXONE SL as permitted according to this supplemental label, read and follow all applicable directions, restrictions, and precautions on the EPA registered label on or attached to the pesticide product container. This Supplemental Labeling contains revised use instructions and or restrictions that may be different from those that appear on the container label. This Supplemental Labeling must be in the possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**syngenta**

## DIRECTIONS FOR USE

Crop	Maximum Number of Applications	Use Pattern	Gramoxone Inteon Rate/Acre	Minimum Total Spray per Acre	Grazing or Preharvest Interval	Precautions, Restrictions and Comments
<b>PEANUTS</b>  For Use Only in the States of Florida and Georgia	2	Broadcast Prior to Planting	1.9-3.75 pts.	Ground: 10 gals.	-	<ul style="list-style-type: none"> <li>For burndown of weeds in peanut fields prior to planting.</li> <li>Use the higher rate on dense populations and/or on larger or harder to control weeds. Apply when weeds are actively growing and between 1" to 6" in height. Vegetation 6" or taller may not be controlled.</li> <li>Gramoxone Inteon may be tank-mixed with other herbicides registered for this use for improved weed control.</li> <li>Always add either: 1) Nonionic surfactant containing 75% or more surface agent at 0.25% v/v (2 pints per 100 gallons), of the finished spray volume, <b>OR</b> 2) Crop oil concentrate (non phytotoxic) containing 15 to 20% approved emulsifier, at 1% v/v (1 gallon per 100 gallons) of the finished spray volume).</li> <li>Weeds emerging after application will not be controlled.</li> <li>DO NOT apply by air.</li> </ul>

Gramoxone® SL and the Syngenta logo trademark of a Syngenta Group Company

©2011 Syngenta

GRAMOXONE SL 1217B-S2 0711 – tb – 08/24/11