

Metalosate®



Zinc

Liquid Amino Acid Foliar



Net Weight: 26.6 lbs./12.0 kg (2.5 Gallons)

v09/2007



07311-CA-GL0025

Lot #:



0700000

METALOSATE® ZINC

1.9-0-0

GUARANTEED ANALYSIS (w/w):

Total Nitrogen (N)	1.9%
(1.9% water soluble organic nitrogen)	
Zinc (Zn)	6.8%

Derived from a complex of soluble salt with partially hydrolyzed vegetable protein.

GENERAL INFORMATION:

Metalosate® Zinc is designed for foliar application on plants to prevent or correct nutrient deficiencies that may limit crop growth and yields. It is water soluble and nontoxic to plants when applied as directed. For best results, apply **Metalosate® Zinc** according to recommendations based on plant tissue or soil analysis.

DIRECTIONS:

Shake well before use. Keep from freezing.

Dilute one part **Metalosate® Zinc** with 20 or more parts water. Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non ionic wetting agent may improve spray coverage of certain hard to wet plants. Do not apply undiluted.

Metalosate® Zinc can be included in a regular spray program on crops. Consult with an **Albion®** representative on compatibility with other spray materials.

The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full sized plants. Reduce the rates proportionately when spraying smaller plants.

Metalosate® Zinc may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

RECOMMENDATIONS:

Field Crops and Vegetables: Apply 8 to 24 fluid ounces per acre during periods of rapid growth or nutritional stress. The application may be repeated 2 or more times through the growing season.

Tree Crops: Make an application of 16 to 48 fluid ounces per acre after the beginning of active growth. The application may be repeated at 2 to 4 week intervals through the growing season. On deciduous trees, 24 to 32 fluid ounces may be applied in the oil sprays at the delayed dormant stage; post harvest applications of 32 to 48 fluid ounces may be made while active, green leaves remain on the trees.

Grapes and Berries: Make an application of 16 to 32 fluid ounces per acre after active growth begins. The application may be repeated at intervals of one week or more through the vegetative growth period.

LIMITED WARRANTY:

Seller's and Manufacturer's obligation is limited to replacement of defective product. **NEITHER SELLER NOR MANUFACTURER SHALL BE LIABLE FOR ANY INJURY, LOSS, OR DAMAGE DIRECTLY OR CONSEQUENTLY ARISING OUT OF THE MISUSE OR INABILITY TO USE THE PRODUCT. ALL OTHER WARRANTIES EXCEPT IN WRITING FROM THE MANUFACTURER, WHETHER EXPRESSED OR IMPLIED ARE HEREBY DISCLAIMED.**

General information regarding the contents and levels of metals in this product is available by calling 1 866 CHELATE (1 866 243 5283).

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>.

Albion, Albion Advanced Nutrition, Metalosate, and the Albion Advanced Nutrition logo are registered trademarks of Albion Laboratories, Inc.

Manufactured by: **Albion Advanced Nutrition®**
101 North Main Street
Clearfield, Utah 84015 U.S.A.

CAv09/2007