



FIFRA

Section 2(ee) Recommendation

Date: July 6, 2016

Product: Agri-Mek® SC Miticide/Insecticide
EPA Reg. No. 100-1351

Use: For Suppression of Broad Mites and Cyclamen Mites in Strawberries

State: Florida

Directions for Use

- **It is a violation of Federal law to use this product in a manner inconsistent with its labeling.**
- **The user must refer to the federally approved labels for the above noted product and read and follow all directions for use, restrictions, and precautions.**
- **The user should have this recommendation in its possession at the time of use.**
- **This recommendation for use of this product is permitted under Section 2(ee) of FIFRA and has not been submitted to or been approved by EPA.**

Agri-Mek SC Miticide/Insecticide will provide suppression of Broad Mite (*Polyphagotarsonemus latus*) and Cyclamen Mite (*Phytonemus pallidus*) when applied at the rate of 3.5 fl oz/A.

To avoid illegal residues Agri-Mek SC Miticide/Insecticide must be mixed with a non-ionic activator type wetting, spreading and/or penetrating spray adjuvant. Do not use binder sticker type adjuvants.

Use sufficient water volume and spray pressure to ensure thorough spray coverage. Direct spray contact is essential to suppressing Broad Mites and Cyclamen Mites.

Spray coverage should be directed into the crown of the plant, specifically newly unfolding leaves and the calyx of newly emerged flower buds. This will enhance the results of an Agri-Mek SC Miticide/Insecticide for suppression of Broad Mites and Cyclamen mites.

Make 2 applications 7-10 days apart when mites first appear.

Wait 21 days after the second application before repeating applications.

Agri-Mek® and the Syngenta logo are trademarks of a Syngenta Group Company

©2016 Syngenta

2(ee) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Expiration Date: July 6, 2020

Label Code: FL1351077AC0716