



Performance

Quality

Value

RESTRICTED USE PESTICIDE

Due to toxicity to Fish and Aquatic Organisms. For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



EPA Reg. No. 34704-978

Product Bulletin

For use on Soybeans to control bean leaf beetles.

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF CONNECTICUT, DELAWARE, ILLINOIS, INDIANA, IOWA, KANSAS, KENTUCKY, MAINE, MARYLAND, MASSACHUSETTS, MINNESOTA, MISSOURI, NEBRASKA, NEW JERSEY, NORTH DAKOTA, OHIO, PENNSYLVANIA, RHODE ISLAND, SOUTH DAKOTA, VERMONT, VIRGINIA, WEST VIRGINIA AND WISCONSIN.

FIFRA Section 2(ee) Recommendation: This recommendation is made and permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U. S. Environmental Protection Agency. This bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

RECOMMENDED APPLICATION		
Use Site	Insect	Rate of TOMBSTONE HELIOS®
Soybeans	Bean Leaf Beetles	1.0 – 1.6 fl. oz./A

Apply specified dosage per acre as a foliar spray when bean leaf beetles begin to migrate into soybean fields. **TOMBSTONE HELIOS®** should be applied when soybeans are between growth stages VC-V2. To control infestations occurring in soybean past the V2 growth stage or in areas where Bean Pod Mottle Virus transmission is a concern, refer to application rate information contained on the **TOMBSTONE HELIOS®** label. A maximum of 2.8 fl. oz./A (0.044 lb. a.i./A) may be applied within a 7 day period. Total applications may not exceed 11.2 fl. oz. of **TOMBSTONE HELIOS®** (0.175 lb. a.i./A) per season.