

## RESTRICTED USE PESTICIDE

Due to ground and surface water concerns. For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's Certification.

Application Type

**AG**

Agricultural



SipcamAdvan

# Stalwart® Xtra Herbicide

## For Weed Control in Corn (All Types)

### ACTIVE INGREDIENT:

Atrazine (2-chloro-4-ethylamino-6-isopropylamino-s-triazine)*	33.0%
Atrazine Related Compounds	0.7%
Metolachlor: 2-chloro- N-(2-ethyl-6-methylphenyl)-N-(2-methoxy-1-methylethyl) acetamide	26.1%

**OTHER INGREDIENTS:** 40.2%

**TOTAL:** 100.0%

This product contains 3.1 lbs. atrazine + related compounds per gallon and 2.4 lbs. of metolachlor active ingredient per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for first aid, directions for use,  
and additional precautionary statements.

**STALWART EXTRA MAY NOT BE USED IN ACCORDANCE WITH  
THE LABEL ON SWEET CORN AFTER DECEMBER 31, 2010**

EPA Reg. No. 60063-23

EPA Est. No. 70815-GA-002 (Lot No. begins with CB)  
EPA Est. No. 70989-MO-1 (Lot No. begins with ST)  
EPA Est. No. 086555-MO-001 (Lot No. begins with AF)

MANUFACTURED FOR:  
**Sipcam Agro USA, Inc.**  
2520 Meridian Parkway, Suite 525  
Durham, NC 27713

9500400-000-20091029

**Net Contents: 2.5 Gallons (9.46 Liters)**

FIRST AID	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have affected person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>Emergency phone numbers</b>	(800) 424-9300 CHEMTREC (transportation and spills) (800) 858-7378 NPIC (human and animal health)
<b>Note to Physician:</b> If ingested, induce emesis or lavage stomach. Administration of an aqueous slurry of activated charcoal can be considered. Treat symptomatically.	

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

# CAUTION:

Harmful if swallowed, inhaled, or absorbed through skin. Do not breathe vapors or spray mist. Avoid contact with eyes, skin, or clothing. Causes moderate eye irritation. Prolonged or frequent skin contact may cause allergic reactions in some individuals.

### PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton. If you want more options, follow the instructions for Category H on an EPA chemical resistance category selection chart.

#### Mixers, loaders, applicators, flaggers, and other handlers must wear:

- Coveralls over Long-sleeved shirt and long pants,
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or viton
- Chemical-resistant footwear plus socks,
- Chemical-resistant headgear (if overhead exposure), and
- A chemical-resistant apron, when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to the concentrate.

See **Engineering Controls** for additional requirements.

**User Safety Requirements:** Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product’s concentrate. Do not reuse them.

## ENGINEERING CONTROLS

Mixers and loaders supporting aerial applications must use a closed system that meets the requirements for dermal protection listed in the Worker Protection Standard (WPS) for Agricultural Pesticides [40 CFR 170.240(d)(4)] and must:

- Wear the long-sleeved shirt and long pants, chemical resistant gloves made of waterproof materials in category H, shoes and socks,
- Wear protective eyewear if the system operates under pressure, and
- Be provided and have immediately available for use in an emergency, such as a spill or equipment breakdown: coveralls, chemical resistant footwear, and chemical resistant apron.

Pilots must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)]. Pilots must wear the PPE required on this labeling for applicators, however, they need not wear chemical-resistant gloves when using an enclosed cockpit.

Flaggers supporting aerial applications must use an enclosed cab that meets the definition on the Worker Protection Standard for Agricultural Pesticides [40 CFR 170.240(d)(5)] for dermal protection. In addition, flaggers must:

- Wear the PPE required on this labeling for flaggers.
- Be provided, have immediately available, and use in an emergency when they must exit the cab in the treated area:
  - Coveralls, chemical resistant gloves made of waterproof materials, chemical resistant footwear, and chemical resistant headgear (if overhead exposure);
- Take off any PPE that was worn in the treated area before reentering the cab;
- Store all such PPE in a chemical resistant container, such as a plastic bag, to prevent contamination of the cab.

When applicators use enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide contains atrazine, which is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water and rinsate.

### Ground Water Advisory

**Stalwart Xtra** contains both the active ingredients atrazine and metolachlor.

Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater.

Metolachlor has the potential to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

### Surface Water Advisory

Metolachlor has the potential to contaminate surface water through ground spray drift. Under some conditions, metolachlor may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow groundwater, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain to surface water.

### Mixing/Loading Instructions

Care must be taken when using this product to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates. Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product must not be mixed or loaded, or used within 50 ft. of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or washwater, and rainwater that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading sites.

Additional State imposed requirements regarding well-head setbacks and operational area containment must be observed.

This product must not be mixed or loaded within 50 ft. of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product must not be applied aerially or by ground within 66 ft. of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 ft. around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66-ft. buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

One of the following restrictions must be used in applying atrazine to tile-outletted fields containing standpipes:

- (1) Do not apply this product within 66 feet of standpipes in tile outletted fields.
- (2) Apply this product to the entire tile-outletted field and immediately incorporate it to a depth of 2-3 inches in the entire field.
- (3) Apply this product to the entire tile outletted field under a no till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW. Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through [[www.atrazine-watershed.info](http://www.atrazine-watershed.info)], or [1-866-365-3014]. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Sipcam Agro USA, Inc. for a refund.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or viton
- Shoes plus socks

### FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

**Note: Not for sale, use, or distribution in Nassau County or Suffolk County, New York.**

## USE INFORMATION

**Stalwart Xtra** is a selective herbicide that may be used before planting, before or after emergence (see directions) for control of most annual grasses and broadleaf weeds in corn (all types). This product may be tank mixed with other herbicides specified on this label for weed control in conventional, minimum-till, and no-till corn.

**Note:** Tank mixtures are permitted only in those states where the tank mix partner is registered. Refer to and follow the label for each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

Following many years of continuous use of atrazine (one of the ingredients in **Stalwart Xtra**), and products chemically related to atrazine, biotypes of some of the weeds listed on this label which are controlled by the atrazine component have been reported to develop resistance to this and chemically related herbicides. Where this is known or suspected, and weeds controlled by this product are expected to be present along with resistant biotypes, we recommend the use of **Stalwart Xtra** in combination or in sequence with registered herbicides, which do not contain triazines. Consult with your State Agricultural Extension Service for specific recommendations.

When tank-mixing or sequentially applying atrazine or products containing atrazine to corn, the total pounds of atrazine applied (lbs a.i./A) must not exceed 2.5 pounds active ingredient per year.

**Stalwart Xtra** alone or in tank mixture with atrazine, isoxaflutole (Balance<sup>®</sup>), metolachlor (Stalwart C), or simazine (Sim-Trol<sup>®</sup>) may be applied early preplant, preplant surface, preplant incorporated, or preemergence in water or fluid fertilizer. Apply post-emergence treatments of **Stalwart Xtra** using water only as the carrier. **Stalwart Xtra** may be applied in tank mix combination with paraquat (Gramoxone<sup>®</sup> Xtra), 2,4-D + glyphosate (Landmaster<sup>®</sup> BW), or glyphosate (Roundup<sup>®</sup>) with or without the above herbicides preplant surface or preemergence.

**Stalwart Xtra** may be applied in water by aircraft. Applications in fluid fertilizer should be only by ground equipment.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

Do not apply this product through any type of irrigation system.

Do not apply this product in a greenhouse.

Do not graze or feed forage from treated areas for 60-days following application for corn (all types). (60-day PHI)

Post-emergent applications to corn must be made before the crop reaches 12 inches in height.

Do not apply under conditions, which favor runoff or wind erosion of soil containing this product to nontarget areas.

To prevent off-site movement due to runoff or wind erosion:

1. Avoid treating powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
2. Do not apply to impervious substrates, such as paved or highly compacted surfaces.
3. Do not use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

Dry weather following preemergence application of **Stalwart Xtra** or a tank mixture may reduce effectiveness. Cultivate if weeds develop.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with other materials, or crop damage or sprayer clogging of the application device may occur.

## MIXING PROCEDURES

Shake 2.5 gal. jugs well or thoroughly recirculate larger containers and bulk tanks before using. **Stalwart Xtra** is a liquid that may be mixed with water or fluid fertilizer and applied as a spray. **Stalwart Xtra** may also be sprayed onto dry bulk granular fertilizer and applied with the granular fertilizer.

### Dry Bulk Granular Fertilizers

Many dry bulk granular fertilizers may be impregnated or coated with **Stalwart Xtra** and used to control weeds in corn.

When applying **Stalwart Xtra** with dry bulk granular fertilizer, follow all directions for use and precautions on the **Stalwart Xtra** label regarding target crops, rates per acre, soil texture, application methods (including timing of application), and rotational crops.

All individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application are the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Impregnation of bulk fertilizer is restricted to commercial facilities. On-farm fertilizer impregnation is prohibited. No more than 500 tons of dry bulk fertilizer can be impregnated per day. No single facility may impregnate fertilizer with this product for more than 30 days per calendar year. The commercial facility impregnating the dry bulk fertilizer must inform, in writing, the user (applicator) of the dry bulk fertilizer that:

- Applicators must wear long-sleeved shirt, long pants, shoes, and socks.
- The restricted-entry interval is 24 hours.

Prepare the herbicide/fertilizer mixture by using any closed drum, belt, ribbon, or other commonly used dry bulk fertilizer blender. Nozzles used to spray **Stalwart Xtra** onto the fertilizer must be placed to provide uniform spray coverage. Care should be taken to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material, such as Agsor<sup>®</sup> or Celatom MP-79<sup>®</sup>, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Absorptive materials should be added only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application result will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Calculate the amount of **Stalwart Xtra** to be used by the following formula:

$$\begin{array}{rcccl} \frac{2,000}{\text{lbs. of fertilizer per acre}} & \times & \text{pts./A of liquid or flowable product} & = & \text{pts. of liquid or flowable product per ton of fertilizer} \\ \\ \frac{2,000}{\text{lbs. of fertilizer per acre}} & \times & \text{lbs./A of dry product} & = & \text{lbs. of dry product per ton of fertilizer} \end{array}$$

### Pneumatic (Compressed Air) Application

High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixtures to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix **Stalwart Xtra** with Exxon Aromatic 200 at a rate of 2.0-2.5 pts./gal. of **Stalwart Xtra**. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Drying agents should not be used when using Aromatic 200.

**Notes:** (1) Mixtures of **Stalwart Xtra** and Aromatic 200 must be used on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications. (2) When impregnating **Stalwart Xtra** in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size is recommended. (3) Drying agents are not recommended for use with On-The-Go impregnation equipment.

**Precautions:** To avoid potential for explosion, (1) Do not impregnate **Stalwart Xtra** on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers. (2) Do not combine **Stalwart Xtra** with a single superphosphate (0-20-0) or treble superphosphate (0-46-0). (3) Do not use **Stalwart Xtra** on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.

### Application

Apply 200-700 lbs. of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential in order to prevent possible crop injury or injury to subsequent rotational crops. Nonuniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil is recommended to obtain satisfactory weed control. On fine- or medium-textured soils in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On coarse-textured soils, make applications approximately 14 days prior to planting.

**Precautions:** (1) To help avoid rotational crop injury, make applications as early as possible, since **Stalwart Xtra** impregnated onto dry bulk granular fertilizers can be expected to last longer in the soil than when **Stalwart Xtra** is applied as a spray in water or fluid fertilizer. (2) To avoid potential crop injury, do not use the herbicide/fertilizer mixture on crops where planting beds are to be formed.

### Application in Water or Fluid Fertilizers

**Stalwart Xtra Alone:** Fill the spray tank 1/2-3/4 full with water or fluid fertilizer, add the proper amount of **Stalwart Xtra**, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform suspension.

**Tank Mixtures:** Fill the spray tank 1/2-3/4 full with water or fluid fertilizer, add the proper amount of **Stalwart Xtra**, then add atrazine, dicamba, isoxaflutole (Balance), linuron, or simazine; next add metolachlor; then add paraquat, 2,4-D + glyphosate (Landmaster BVV), or glyphosate, depending on the tank mix combination desired; and finally, add the rest of the water or fluid fertilizer. Only water may be used with **Stalwart Xtra** + Liberty<sup>®</sup> Herbicide when applied postemergence to corn designated as tolerant to Liberty (glufosinate); and with Roundup Ultra<sup>®</sup> when applied postemergence to corn designated as tolerant to Roundup (glyphosate). Provide sufficient agitation during mixing and application to maintain a uniform suspension.

### Compatibility Test

A jar test is recommended before tank mixing to ensure compatibility of **Stalwart Xtra** with other pesticides. The following test assumes a spray volume of 25 gals./A. For other spray volumes, make appropriate changes in the ingredients.

**Note:** Nitrogen solutions or complete liquid fertilizers may replace all or part of the water in the spray. Since liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) each time before use.** Incompatibility of tank mixtures is more common with suspensions of fertilizer and pesticides. Commercial application equipment may improve compatibility in some instances. Check compatibility using this procedure:

1. Add 1.0 pt. of carrier (fertilizer or water) to each of 2 one-qt. jars with tight lids. **Note:** Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
2. To **one** of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent approved for this use (1/4 tsp. is equivalent to 2.0 pts./100 gals. spray). Shake or stir gently to mix. When an adjuvant is to be used with this product, Sipcam Agro USA recommends the use of Complex®, Unite® or a Chemical Producers and Distributors Association (CPDA) certified adjuvant.
3. To **both** jars, add the appropriate amount of pesticide(s) in their relative proportions based on specified label rates. If more than one pesticide is used, add them separately with dry herbicides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicides for this test follows:  
**Dry herbicides:** For each pound to be applied per acre, add 1.5 level teaspoons to each jar.  
**Liquid herbicides:** For each pint to be applied per acre, add 0.5 teaspoon or 2.5 milliliters to each jar.
4. After adding all ingredients, put lids on and tighten, and invert each jar 10 times to mix. Let the mixtures stand 15-30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the 2 jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (A) slurry the dry pesticide(s) in water before addition, or (B) add 1/2 of the compatibility agent to the fertilizer or water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.
5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **Storage and Disposal** section in this label.

## SOIL TEXTURE INFORMATION

Within rate ranges in all tables on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

**Directions are based upon soil textural classes, generally categorized as follows:**

Coarse	Medium	Fine
Sand	Loam	Sandy clay loam
Loamy sand	Silt loam	Silty clay loam
Sandy loam	Silt	Clay loam
		Sandy clay
		Silty clay
		Clay

## APPLICATION INFORMATION

**Ground Application:** Use sprayers that provide accurate and uniform application. Screens in nozzles and in suction and in-line strainers should be no finer than 50-mesh. Use a pump with capacity to: (1) maintain 35-40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Unless otherwise specified, use a minimum of 10 gals. of spray mixture per acre. Rinse sprayer thoroughly with clean water immediately after use.

For band applications, calculate amount to be applied per acre as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

**Low Carrier Application (Broadcast Ground Application Only):** Use sprayers, such as Ag-Chem RoGator®, Hagie, John Deere Hi-Cycle™, John Deere 4700 Sprayer, Melroe Spra-Coupe, Tyler Patriot™, or Willmar Air Ride®, that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35-40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5.0 gals. of spray mixture per acre. Maximum recommended sprayer speed is 15 mph. Maintain uniform travel speed while spraying. Rinse sprayer thoroughly with clean water immediately after each use.

**Note:** Low pressure nozzles are recommended to reduce drift and increase application accuracy. Care should be taken when using automatic rate-controlling devices to spray the material within the rated working pressure and flow ranges of the nozzle selected. Nozzle screens should be used when recommended by the manufacturer. All nozzles should be placed on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended. Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

**Aerial Application (for Stalwart Xtra alone):** Use aerial application only where broadcast applications are specified. Apply a minimum of 1.0 gal. of water for each 1.0 gal. of this product applied per acre, but for rates below 1.0 gal./A, use in sufficient water to equal 2.0 gals./A of total spray. Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply **Stalwart Xtra** by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders should avoid inhalation of spray mist and prolonged contact with skin.

### Aerial Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment- and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outermost nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downward more than 45 degrees.

Where states have more stringent regulations, they must be observed.

The applicator should be familiar with and take into account the information covered in the **Aerial Drift Reduction Advisory Information** section below.

## **Aerial Drift Reduction Advisory Information**

### **Information on Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see **Wind, Temperature and Humidity**, and **Temperature Inversions**).

### **Controlling Droplet Size**

- **Volume** – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- **Pressure** – Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.
- **Number of Nozzles** – Use the minimum number of nozzles that provide uniform coverage.
- **Nozzle Orientation** – Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- **Nozzle Type** – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

### **Boom Length**

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

### **Application Height**

Applications must not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

### **Swath Adjustment**

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

### **Wind**

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### **Temperature and Humidity**

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

### **Temperature Inversions**

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### **Sensitive Areas**

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

### **Rate Limitations**

Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or greater setbacks), which are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

**Note:** For purposes of calculating total atrazine active ingredient applied, **Stalwart Xtra** contains 3.1 lbs. a.i. atrazine + related compounds per gal. (0.775 lb. a.i./qt.).

#### **For All Soil Applications**

Apply a maximum of 2.0 lbs. a.i./A (2.6 qts. of product/A) as a single preemergent broadcast application on soils that are not highly erodible or on highly erodible soils (as defined by the Natural Resources Conservation Service) if practicing conservation tillage and at least 30% of the soil surface is covered with plant residues at planting; or apply a maximum of 1.6 lbs. a.i./A (2.1 qts. of product/A) as a single preemergent broadcast application on highly erodible soils (as defined by the Natural Resources Conservation Service) if less than 30% of the soil surface is covered with plant residues at planting; or 2.0 lbs. a.i./A (2.6 qts. of product/A) if only applied postemergent.

#### **FOR POSTEMERGENCE APPLICATION TO CORN**

If no atrazine was applied prior to corn emergence, apply a maximum of 2 lbs. a.i./A (2.6 qts./A) broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. a.i./A (3.2 qts. of **Stalwart Xtra**) per calendar year.

## ROTATIONAL CROPS

Do not rotate to food or feed crops other than those listed below:

(1) If treated crop is lost due to poor germination, hail, flood, insects, etc., corn may be replanted immediately. Do not make a second broadcast application. If the original application was banded and the second crop is planted in the untreated row middles, a second banded treatment may be applied. (2) Corn, sorghum, soybeans, cotton, or peanuts may be planted the spring following treatment. Do not graze or feed forage or fodder from cotton to livestock, or illegal residues may result. (3) Injury may occur to soybeans planted the year following application on soils having a calcareous surface layer. (4) In eastern parts of the Dakotas, KS, western MN, and NE, do not rotate to soybeans for 18 months following application if the rate applied to corn or sorghum was more than 2.0 lbs. a.i. of atrazine or equivalent band application rate, or soybean injury may occur. (5) If applied after June 10, do not rotate with crops other than corn or sorghum the next year, or crop injury may occur. (6) In the High Plains and Inter-mountain areas of the West, where rainfall is sparse and erratic or where irrigation is required, use only when corn or sorghum is to follow corn or sorghum, or a crop of untreated corn or sorghum is to precede other rotational crops. (7) Do not plant sugar beets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes the year following application, or injury may occur.

## WEED CONTROL INFORMATION

**Stalwart Xtra** is effective for the control of grassy and broadleaf weed plants that can infest agricultural fields. The spectrum of effective weed control may differ depending upon the application pattern and timing of application(s) of this product. The following table is a partial illustration of the potential usefulness of this product to mitigate the growth of weeds.

Weeds Controlled – Preplant and Preemergence Applications			
barnyardgrass (watergrass)	green foxtail	carpetweed	jimsonweed
browntop panicum	prairie cupgrass	chickweed	lambsquarters
crabgrass	red rice	cocklebur*	morningglory
crowfootgrass	signalgrass ( <i>Brachiaria</i> )*	common ragweed	mustards
fall panicum	southwestern cupgrass	Florida pusley	nightshades
foxtail millet	witchgrass	galinsoga	pigweed
giant foxtail	yellow foxtail	giant ragweed*	smartweed
goosegrass	yellow nutsedge*	henbit	velvetleaf*
			waterhemp
Weeds Controlled – Postemergence Applications			
barnyardgrass (watergrass)	fall panicum	jimsonweed	prickly sida
cocklebur	flixweed	kochia	purslane
common ragweed	giant foxtail	lambsquarters	ragweed
crabgrass	green foxtail	morningglory	smartweed
crowfootgrass	yellow foxtail	mustard	velvetleaf
		pigweed	waterhemp
Weeds Partially Controlled**			
sandbur	shattercane	volunteer sorghum	yellow nutsedge
seedling johnsongrass	sicklepod	woolly cupgrass	common purslane

\* Control of these weeds can be erratic, especially under dry weather conditions. Control escaped weeds with cultivation or application of an appropriate EPA-registered postemergence herbicide. On fine-textured soils, only partial control can be expected.

\*\* Control may be improved by following these suggested procedures:

1. Apply up to the maximum single application rate in Table 1 for your given soil texture and rate limitation based on your soil conservation practices.
2. **Thoroughly till moist soil** to destroy germinating and emerged weeds. If **Stalwart Xtra** is to be applied preplant incorporated, this tillage may be used to incorporate **Stalwart Xtra** if uniform 2-inch incorporation is achieved as recommended under **Application Procedures**.
3. Plant crop into moist soil **immediately after tillage**. If **Stalwart Xtra** is to be used preemergence, apply at planting or immediately after planting.
4. If available, **sprinkler irrigate** within 2 days after application. Apply 1/2-1 inch of water. Use lower water volume (1/2 inch) on coarse-textured soils and higher volume (1 inch) on fine-textured soils.
5. If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.



# APPLICATION PROCEDURES

## Early Preplant

If you apply this product early preplant, use the appropriate rate from Table 1.

Table 1: Stalwart Xtra – Early Preplant Rates of Application			
Soil Texture	Single Application	Split Application*	
		30-45 Days Pre-planting	At Planting
<b>COARSE</b> Sand, loamy sand, sandy loam	2.1 qts./A	DO NOT APPLY	
<b>MEDIUM</b> Loam, silt loam, silt	<b>A.</b> 2.1 qts./A	1.4 qts./A	0.7 qt./A
	<b>B.</b> 2.1-2.6 qts./A	1.4 qts./A to 1.75 qts./A	0.7 qt./A to 0.9 qt./A
<b>FINE</b> Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	<b>A.</b> 2.1 qts./A	1.4 qts./A	0.7 qt./A
	<b>B.</b> 2.6 qts./A	1.75 qts./A	0.9 qt./A

\* Split applications can be made less than 30 days before planting if desired.

**A.** Do not exceed this rate on highly erodible land with less than 30% plant residue cover. Control of certain weeds may be reduced and a tank mix partner or an application of a postemergence herbicide may be needed.

**B.** Do not exceed this rate on soils that are not highly erodible or on highly erodible soils if at least 30% of the soil is covered with plant residue.

Applications made less than 30 days prior to planting may be as either a split or single treatment. Use the lower rate for light expected weed infestations and the higher rate for heavy expected weed infestations. On *coarse-textured soils*, apply 2.1 qts./A not more than 2 weeks prior to planting. The above procedure may be followed if atrazine or metolachlor or simazine is used in tank mixtures with **Stalwart Xtra**. Substitute a fluid fertilizer for some or all of the water carrier for burndown of existing annual weeds listed on this label up to the 2-leaf stage of development. The addition of crop oil concentrate to the spray mixture will enhance the burndown activity. If larger weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, paraquat or glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

Use on *medium- and fine-textured soils* with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WY.

On *medium- and fine-textured soils* with minimum- or no-tillage systems in DE, MD, MI, NY, OH, PA, VA, and WV, early preplant applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., atrazine, primisulfuron-methyl (Beacon®), dicamba (Banvel®), bentazone (Basagran®), bromoxynil (Brominal® or Buctril®), primisulfuron-methyl + prosulfuron (Exceed®), atrazine + dicamba (Marksman®), or 2,4-D. If the postemergence treatment includes the herbicide used early preplant, do not exceed the labeled rate for corn on a given soil texture. Observe all directions for use, precautions, and limitations on the label of the post-emergent herbicide.

**Stalwart Xtra** may be used according to the above directions to control winter wheat planted as a cover crop in IN, KY, and OH, in addition to providing residual weed control. The wheat must be less than 6 inches tall (preferably still in a dormant or semi-dormant state coming out of winter) at the time of application. Depending on rainfall, 10-20 days may be required to completely kill the wheat. In the event that adequate rainfall does not occur, control of the winter wheat may be unsatisfactory and the application of a contact herbicide (i.e., paraquat or glyphosate) may be required before planting the crop.

On *medium- and fine-textured soils* following final seedbed preparation in the Blacklands and Gulf Coast areas of TX, an early preplant application of **Stalwart Xtra** at 1.6-1.9 qts./A may be made 30-45 days before planting. Grass suppression of 2-3 weeks after planting can be expected as a result of this application. Do not incorporate or disturb the soil before planting, and avoid moving the soil during the planting operation. A follow-up application of metolachlor may be needed in fields with a history of heavy grass pressure. Apply after planting, but before corn and grass weeds emerge.

**Notes:** (1) If a follow-up application of metolachlor is needed, do not exceed a total of 1.6 lbs. a.i. of metolachlor per acre, including the preplant **Stalwart Xtra** application on *medium- or fine-textured soils*. On *fine-textured soils* with more than 3% organic matter, do not exceed 1.9 lbs. a.i. of metolachlor.

[To determine the total lbs. a.i. of metolachlor per acre, use the following 2-step method:

A. Determine the lbs. of metolachlor applied as **Stalwart Xtra** (1 qt = 0.6 lb. a.i. of metolachlor); then,

B. If metolachlor is to be used, add the lbs. a.i. to be applied in these products to the lbs. in Step A above.]

(2) To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

## Preplant Surface, Preplant Incorporated, or Preemergence

If you apply this product preplant surface, preplant incorporated, or preemergence, use the appropriate rate from Table 2.

**Preplant Surface:** Apply uniformly to the soil surface within 14 days before planting. Where applications are made to coarse soils more than 7 days before planting, use the rates in Table 1.

**Preplant Incorporated:** Apply to the soil and incorporate into the top 2 inches of the soil within 14 days before planting, using a finishing disk, finishing harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use the preplant incorporated method if furrow irrigation is used or when a period of dry weather after application is expected. If crop is to be planted on beds, apply and incorporate after bed formation.

**Preemergence:** Apply to the soil surface at planting (behind the planter) or after planting, but before weeds or crop emerge.

**Table 2: Stalwart Xtra – Preplant Surface, Preplant Incorporated, or Preemergence Rates of Application**

Soil Texture	Broadcast Rate Per Acre	
	Less Than 3% Organic Matter	3% Organic Matter or Greater
<b>COARSE</b> Sand, loamy sand, sandy loam	1.3 qts.	1.6 qts.
<b>MEDIUM</b> Loam, silt loam, silt	1.6 qts.	2.1 qts.
<b>FINE</b> Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	2.1 qts.	<b>A.</b> 2.1 qts.
		<b>B.</b> 2.1-2.6 qts.*
Muck or peat soils (more than 20% organic matter)	DO NOT USE	

\* For cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter: Apply 2.6 qts. of **Stalwart Xtra** per acre.

**A.** Do not exceed this rate on highly erodible land with less than 30% plant residue cover. Control of certain weeds may be reduced and a tank mix partner or an application of a postemergence herbicide may be needed.

**B.** Do not exceed this rate on soils that are not highly erodible or on highly erodible soils if at least 30% of the soil is covered with plant residue.

**Notes:** (1) In the event of escape of annual weeds following an early preplant, preplant surface, preplant incorporated, or preemergence treatment of **Stalwart Xtra** applied alone or in combination, follow with a post-emergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., atrazine, primisulfuron-methyl (Beacon), nicosulfuron (Accent<sup>®</sup>), dicamba (Banvel), bentazone (Basagran), bromoxynil (Brominal or Buctril), primisulfuron-methyl + prosulfuron (Exceed), atrazine + dicamba (Marksman), or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, do not exceed the labeled rate for corn on a given soil texture. (2) Bromoxynil may be applied postemergence alone or in tank mix combination with atrazine. Do not exceed 1.2 lbs. a.i./A of atrazine in tank mix combination with bromoxynil postemergence. Refer to the atrazine and bromoxynil labels for specific rates and precautions. (3) If atrazine or another product containing atrazine is used post-emergence following application of **Stalwart Xtra**, do not exceed a total of 2.5 lbs. a.i./A of atrazine per year. (4) Substitute a fluid fertilizer for some or all of the water carrier for burndown of existing annual weeds listed on this label up to the 2-leaf stage of development. The addition of crop oil concentrate to the spray mixture will enhance the burndown activity. If larger weeds are present, add a contact herbicide as noted in the **Stalwart Xtra Combinations** section of this label.

#### **Postemergence Broadcast**

If you apply this product early postemergence, use the appropriate rate from Table 3. Apply post-emergence treatments of **Stalwart Xtra** using water only as the carrier. Apply this treatment before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control. Occasional corn leaf burn may result, but this should not affect later growth or yield. Do not apply postemergence in fluid fertilizer, or severe crop injury may occur.

**Note:** To avoid possible illegal residues, do not graze or feed forage from treated areas for 60 days following application.

**Table 3: Postemergence Broadcast Rates of Application**

Soil Texture	Broadcast Rate Per Acre
<b>COARSE</b> Sand, loamy sand, sandy loam	1.6 qts.
<b>MEDIUM</b> Loam, silt loam, silt	2.1 qts.
<b>FINE</b> Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	2.1-2.6 qts.*

\* For better residual control of cocklebur, velvetleaf, and yellow nutsedge on *fine-textured soils* above 3% organic matter, apply 2.6 qts. of **Stalwart Xtra** per acre.

**Notes:** (1) If no atrazine was applied prior to corn emergence, apply a maximum of 2 lbs. a.i./A broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. a.i./A per calendar year. (2) If atrazine or atrazine plus metolachlor tank mixtures have been applied early preplant, preplant surface, preplant incorporated, or pre-emergence, limit the **Stalwart Xtra** early post application not to exceed a total of 2.5 lbs. of active ingredient in atrazine or 3.75 lbs. of the active ingredient metolachlor per acre, or illegal residues may result.

#### **Postemergence-Directed**

**Stalwart Xtra** may be applied at 1.3-2.6 qts./A in a minimum of 15 gals. of water as a postemergence-directed treatment to corn to extend control of weeds. Apply using the appropriate rate from Table 4.

For best results, apply **Stalwart Xtra** to weed-free soil following use of a preplant surface, preplant incorporated, or preemergence herbicide, or following a lay-by cultivation. If weeds have emerged at the time of **Stalwart Xtra** application, apply before grass and broadleaf weeds exceed the 2-leaf stage. Application to weeds larger than the 2-leaf stage will generally give unsatisfactory control. Apply to corn not exceeding 12 inches in height. Minimize contact with corn leaves. Do not apply postemergence in fluid fertilizer, or severe crop injury may occur.

**Note:** To avoid possible illegal residues, do not graze or feed forage from treated areas for 60 days following application.

Table 4: Postemergence-Directed Rates of Application	
Soil Texture	Broadcast Rate Per Acre
<b>COARSE</b> Sand, loamy sand, sandy loam	1.3 qts.
<b>MEDIUM</b> Loam, silt loam, silt	2.1 qts.
<b>FINE</b> Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	2.1-2.6 qts.*

\* For better residual control of cocklebur, velvetleaf, and yellow nutsedge on fine-textured soils above 3% organic matter, apply 2.6 qts. of **Stalwart Xtra** per acre.

**Notes:** (1) If no atrazine was applied prior to corn emergence, apply a maximum of 2 lbs. a.i./A broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. a.i./A per calendar year. (2) If atrazine plus metolachlor tank mixtures have been applied preplant surface, preplant incorporated, or preemergence, limit the **Stalwart Xtra** post-directed application not to exceed a total of 2.5 lbs. of the active ingredient atrazine or 3.75 lbs. of the active ingredient metolachlor per acre on a corn crop, or illegal residues may result.

## **STALWART EXTRA COMBINATIONS**

Always follow label instructions for tank mix products when mixing with **Stalwart Xtra**.

\* When tank mixing **Stalwart Xtra** with atrazine formulations, refer to the **Stalwart Xtra Rate Limitations** section of this label. Do not exceed the following:

On highly erodible land with less than 30% plant residue cover prior to crop emergence	1.6 lbs. a.i. of atrazine
On other land prior to crop emergence	2.0 lbs. a.i. of atrazine
Postemergence applications only – any land	2.0 lbs. a.i. of atrazine
Preemergence + postemergence applications	2.5 lbs. a.i. of atrazine

### **I. Tank Mixture with Atrazine, Metolachlor (Stalwart C), Simazine (Sim-Trol), or Isoxaflutole (Balance) – Conventional Tillage**

**Note:** Check the compatibility of **Stalwart Xtra** tank mixtures with isoxaflutole before mixing in spray tank by using the procedure described under **Application in Water or Fluid Fertilizers**.

**Atrazine (4L or 90DF):** Add up to 1.0 qt. of Sipcarn Atrazine 4L (1.1 lbs. of 90DF) per acre to the rate of **Stalwart Xtra** specified in Table 3 in the southeastern U.S. where high rainfall can shorten the duration of control of broadleaf weeds, and in all areas where heavy infestations of cocklebur, morningglory, velvetleaf, or other broadleaf weeds claimed are expected.

**Metolachlor Products:** Add up to 0.33 pt. of **Stalwart C** per acre (0.32 lb. a.i. metolachlor) to the rate of **Stalwart Xtra** specified in Table 3 when heavy infestations of yellow nutsedge, sandbur, or seedling johnsongrass are expected.

**Simazine (4L or 90DF):** Add up to 1.0 qt. of Sim-Trol 4L (1.1 lbs. of Sim-Trol 90DF) per acre to the rate of **Stalwart Xtra** specified in Table 3 where heavy infestations of crabgrass or fall panicum are expected or additional control of certain broadleaves is desired.

**Isoxaflutole (Balance) (Field Corn Only):** The tank mixture of **Stalwart Xtra** + isoxaflutole provides control of weeds listed on the **Stalwart Xtra** label, certain weed biotypes resistant to ALS-inhibitor herbicides and to triazine herbicides, velvetleaf, and others on the respective product labels. Isoxaflutole will contribute to the control of problem grass and other broadleaf species on its label. Application may be preplant (surface-applied up to 14 days before planting), preplant incorporated, or preemergence in conventional tillage, conservation tillage, and no-till systems. Refer to **Table 1: Stalwart Xtra – Early Preplant** for the early preplant application rate (8-14 days before planting) or refer to **Table 2** for the appropriate rate for preplant (surface-applied 0-7 days before planting), preplant incorporated, or preemergence application. Refer to the **Application Procedures and Tank Mix Directions** on the isoxaflutole label, but to reduce the potential for injury from isoxaflutole contact with corn, use 1.0 oz./A of isoxaflutole on coarsely-textured soils and 1.0-1.5 oz./A on medium- and fine-textured soils in conventional, conservation, and no-tillage systems. For early preplant applications 8-14 days before planting, add 0.5 oz./A of isoxaflutole to the rates of isoxaflutole described above.

Observe all applicable directions, precautions, and limitations on the **Stalwart Xtra** and isoxaflutole labels when applying these products in tank mix combination in states where isoxaflutole is registered. Where difficult species and/or severe weed populations are expected, use the maximum rates of **Stalwart Xtra** and isoxaflutole where rate ranges are listed for the tank mixture.

### **2. Tank Mixture of Stalwart Xtra Alone or Stalwart Xtra + Atrazine, Isoxaflutole (Balance), Metolachlor (Stalwart C), or Simazine (Sim-Trol), with Paraquat (Gramoxone Inteon), 2,4-D + Glyphosate (Landmaster BW), or Glyphosate (Roundup) for Minimum-Tillage or No-Tillage Systems**

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, 2,4-D + glyphosate, or glyphosate should be tank mixed with **Stalwart Xtra** alone or with **Stalwart Xtra** + atrazine, isoxaflutole, metolachlor, or simazine. When used as directed, the paraquat portion of the tank mixture controls most emerged annual weeds and suppresses many perennial weeds. 2,4-D + glyphosate or glyphosate combinations will control emerged annual and perennial weeds when applied as directed on its label. The **Stalwart Xtra** portion of the tank mixture provides preemergence control of the weeds listed on this label in the **Stalwart Xtra Alone** section for corn. The addition of atrazine, isoxaflutole, metolachlor, or simazine offers the advantage indicated for each under **Tank Mixture with Atrazine, Metolachlor (Stalwart C), Simazine (Sim-Trol), or Isoxaflutole (Balance) – Conventional Tillage** section.

**Application:** Apply before, during, or after planting, but before corn emerges, at the appropriate rate in Table 5. Up to 1.0 qt. of Sipsam Atrazine 4L (1.1 lbs. of 90DF), or 1.0-2.0 oz. of isoxaflutole (refer to **Tank Mixture with Balance** for specific rate), or 0.33 pt. of metolachlor (Stalwart C), or 1.0 qt. of Sim-Trol 4L (1.1 lbs. of Sim-Trol 90DF) per acre may be added to the rate of **Stalwart Xtra** specified in Table 5. Add paraquat, 2,4-D + glyphosate, or glyphosate at labeled rates. **Tank mixtures with isoxaflutole (Balance) can only be used on field corn.**

Apply in 20-60 gals. of water per acre with conventional spray equipment.

**3. Tank Mixture of Stalwart Xtra Alone or Stalwart Xtra + Atrazine, or Isoxaflutole (Balance), with 2,4-D or 2,4-D + Dicamba (Banvel) for Minimum-Tillage or No-Tillage Systems**

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, **Stalwart Xtra** may be applied in combination with atrazine or isoxaflutole. When used as directed, the **Stalwart Xtra** portion of the tank mixture provides preemergence control of the weeds listed on this label in the **Stalwart Xtra Alone** section for corn. The addition of atrazine or isoxaflutole offers the advantage indicated for each under **Tank Mixture with Atrazine, Metolachlor, Simazine, or Isoxaflutole (Balance) – Conventional Tillage** section.

**Application:** Apply **Stalwart Xtra** before, during, or after planting, but before corn emerges, at the appropriate rate in Table 5. Up to 1.0 qt. of Sipsam Atrazine 4L (1.1 lbs. of 90DF) or 1.0-2.0 oz. of isoxaflutole (refer to **Tank Mixture with Isoxaflutole (Balance)** for specific rate), per acre may be added to the rate of **Stalwart Xtra** specified in Table 5.

Where heavy crop residues exist, add an appropriately labeled 2,4-D amine or low volatile ester to the spray tank last and apply in a minimum of 25 gals. of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore are recommended instead of water. Add X-77® surfactant at 1.0-2.0 qts./100 gals. of diluted spray, or another surfactant cleared for use on growing crops at its recommended rate. Apply before weeds exceed 3 inches in height. If alfalfa is present, add dicamba (Banvel) to the spray mixture at 0.33-0.5 pt./A and apply before alfalfa exceeds 6 inches in height.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add paraquat (Gramoxone Inteon) at the rate of 2.5 pts./A in place of, or in addition to, 2,4-D, as indicated above. Do not apply paraquat in suspension-type liquid fertilizer (using clay suspending agents), as the activity of paraquat will be reduced. Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination.

Table 5: Stalwart Xtra for Minimum-Tillage or No-Tillage Corn	
Soil Texture	Broadcast Rate Per Acre
<b>COARSE</b> Sand, loamy sand, sandy loam	1.6 qts.
<b>MEDIUM</b> Loam, silt loam, silt	2.1 qts.
<b>FINE</b> Sandy clay loam, silty clay loam, clay loam, sandy clay, silty clay, clay	<b>A.</b> 2.1 qts.
	<b>B.</b> 2.1-2.6 qts.*
Muck or peat soils	DO NOT USE

\* For cocklebur, yellow nutsedge, and velvetleaf control on fine-textured soils above 3% organic matter: Apply 2.6 qts. of **Stalwart Xtra** per acre.

**A.** Do not exceed this rate on highly erodible land with less than 30% plant residue cover. Control of certain weeds may be reduced and a tank mix partner or an application of a postemergence herbicide may be needed.

**B.** Use this rate for all other applications.

**4. Tank Mixture with Linuron (Lorox) for Control of Lambsquarters and Pigweed**

For prolonged control of lambsquarters and pigweed in DE, MD, NJ, NY, PA, VA, and WV, **Stalwart Xtra** may be applied preemergence in tank mix combination with linuron. Apply **Stalwart Xtra** according to the rates in Table 2 and linuron according to the following rates.

Soil Texture	Broadcast Rate Per Acre
Sandy loam (1-3% organic matter)	0.67 lb. Lorox*
Sandy loam (3-6% organic matter)	1.0 lb. Lorox*
Medium- and fine-textured soils (1-6% organic matter)	1.0 lb. Lorox*

\* or equivalent linuron formulation, one pt. of Lorox L equals 1.0 lb. of Lorox DF.

Follow instructions and precautions on the **Stalwart Xtra** and linuron labels when tank mixing these products.

**Rotational Crops:** Follow the crop rotation instructions in the **Stalwart Xtra Alone** section for corn.

**5. Tank Mixture with Mesotrione (Callisto®)**

For preemergence control of weeds in corn, **Stalwart Xtra** may be applied in combination with mesotrione. Apply **Stalwart Xtra** according to the rates in Table 2 and mesotrione at a rate of 5.0-6.0 fl. oz./A. Observe all directions for use, precautions, and limitations on the respective product labels when applying these products in tank mix combination.

## 6. Tank Mixtures For Postemergence Weed Control in Field Corn Only

For postemergence control of weeds in specific types of field corn, the **Stalwart Xtra** combinations listed below may be used. Full season weed control from early preplant, preplant incorporated, or preemergence treatments can lead to maximum yield potential under competition-free conditions. However, if control of emerged weeds is needed, a postemergence program listed below can be applied to provide residual control for the remainder of the season.

**Notes:** (1) Follow all label directions, instructions, precautions, and limitations for each product used. (2) Do not use fluid fertilizer with these mixtures or corn injury may occur. (3) For each tank mixture with **Stalwart Xtra**, apply only to the specific field corn type specified on the tank mix product label. (4) In-row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.

- a. **Stalwart Xtra + Liberty Herbicide: Postemergence Use in LibertyLink® Corn or Corn Warranted by Bayer CropScience as Being Tolerant to Liberty Herbicide** - The tank mixture of **Stalwart Xtra** + Liberty Herbicide can be applied postemergence to weeds and corn from seed designated as LibertyLink or corn warranted by Bayer CropScience as being tolerant to Liberty Herbicide. Liberty provides postemergence control of a broad spectrum of grass and broadleaf weeds and the **Stalwart Xtra** provides residual control of grasses and broadleaf weeds listed in the label section **Stalwart Xtra Applied Alone – Weeds Controlled**. For the proper rate of **Stalwart Xtra** applied postemergence, refer to Table 3 and use the minimum rate per soil texture for season-long residual control. Refer to the Liberty label for the Liberty postemergence application rate according to weed species and their maximum height at the time of postemergence application. Where multiple weed species are present, use the highest Liberty rate specified to control the species and growth stages present.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **Stalwart Xtra** and Liberty Herbicide labels.

- b. **Stalwart Xtra + Glyphosate (Roundup Ultra) for Postemergence Application to Corn with the Roundup Ready® Gene** - The tank mixture of **Stalwart Xtra** + Roundup Ultra can be applied postemergence to weeds and to corn designated as containing the Roundup Ready Gene. Application may be applied postemergence to Roundup Ready corn up to 12 inches in height. This mixture will provide postemergence control of weed species on the Roundup Ultra label and also residual control of weed species on the **Stalwart Xtra** label. Use the minimum **Stalwart Xtra** rate postemergence with Roundup Ultra in Roundup Ready corn as specified in Table 3 of this label according to soil texture. Refer to the **Supplemental Labeling of Roundup Ultra for Postemergence Application to Corn with the Roundup Ready Gene** and to each product label and follow all appropriate use directions, application procedures, precautions, and limitations. Apply 24-32 fl. oz./A of Roundup Ultra for control of labeled broadleaf and grass weeds. Refer to the Roundup Ultra label for directions for control of problem species.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the **Stalwart Xtra** and Roundup Ultra labels, and on the **Supplemental Labeling of Roundup Ultra for Postemergence Application to Corn with the Roundup Ready Gene**. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

- c. **Stalwart Xtra + Exceed:** Apply 1.33-1.75 qts./A of **Stalwart Xtra** + 1.0 oz./A of Exceed to corn that is 4-12 inches tall. The application may be broadcast, semi-directed, or directed. The **Stalwart Xtra** rate is based on soil texture, with 1.33 qts./A on *coarse*, and 1.75 qts./A on *medium and fine soils*. Add a nonionic surfactant at 0.25% v/v.

This mixture is effective for control of many annual broadleaf weeds and some grasses. A few instances of broadleaf weed control antagonism have been observed with this combination. Control of certain annual grasses can be improved with the addition of Accent.

- d. **Stalwart Xtra + Exceed + Accent:** Apply the same rates of **Stalwart Xtra** and Exceed as mentioned above. Add Accent at 0.33 oz./A for more effective control of certain annual grasses. Apply to field corn between 4 and 12 inches. Add a nonionic surfactant at 0.25% v/v. The use of fertilizer or crop oil concentrate with this combination may cause injury to corn.

- e. **Stalwart Xtra + Spirit®** - Spirit herbicide at 1.0 oz./A can be substituted in place of Exceed in the above combinations in field corn only.

**Notes:** Do not use fertilizer or crop oil concentrate with these mixtures or injury to field corn may occur. The combination of **Stalwart Xtra** with other products for postemergence weed control in corn is generally not recommended. **These combinations may cause injury and/or weed control concerns that would not exist when the products are used separately.** A certain inherent risk is involved with the various combinations of these products used postemergence in corn. [It should be noted that early preplant, preplant incorporated, or preemergence control of these weeds would usually provide more timely weed control resulting in higher yields than postemergence treatments.]

### Mixing Order

Add these products (Tank Mixtures c, d, and e) to the tank mix in the following order:

1. Products in water-soluble bags should be added first.
2. **Stalwart Xtra**.
3. Additives.

**Precautions:** (1) Follow all label instructions, precautions, and rotational restrictions for individual products when making these applications to field corn. When **Stalwart Xtra** is applied after June 10, crop injury may occur the following year if you rotate to crops other than corn or sorghum. (2) In-row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** This product may be stored at temperatures down to -30°F. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes.

**Pesticide Disposal:** Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office. Open dumping is prohibited.

### Container Disposal:

**Containers < 5 Gallons: Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Minibulk Containers: Non-refillable container.** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Bulk Containers: Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. When the container is empty, replace the cap and seal all openings that have been opened during use; and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call Chem-Trec. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

## NOTICE OF WARRANTY

**Sipcam Agro USA, Inc.** warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of **SIPCAM**. To the extent permitted by law, in no case shall **SIPCAM** be liable for consequential, special or indirect damages resulting from the use or handling of this product. To the extent permitted by law, all such risks shall be assumed by the Buyer. **SIPCAM MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

Stalwart and Sim-Trol are registered trademarks of Sipcam Agro USA, Inc.

Beacon, Callisto, Concep, Exceed, and Gramoxone are registered trademarks of Syngenta Group Company

Accent and Lorox are registered trademarks of E. I. duPont de Nemours and Company, Inc.

Ag-Chem RoGator is a registered trademark of Ag-Chem Equipment Company

Agsorb is a registered trademark of Oil-Dri Corporation

Banvel, BASgran, and Marksman are registered trademarks of BASF Corporation

Celatom MP-79 is a trademark of Eagle-Picher Industries, Inc.

Complex is a registered trademark of KALO Agricultural Chemicals, Inc.

Hi-Cycle™ is a trademark of John Deere Company

Landmaster, Roundup, Roundup Ready, Roundup Ultra and Screen are registered trademarks of Monsanto Company

Balance, Brominal, Butcrlil, Liberty, and LibertyLink are registered trademarks of Bayer CropScience

Tyler Patriot™ is a trademark of Tyler Ltd. Partnership

Unite is a registered trademark of HACO, Inc.

Willmar Air Ride is a registered trademark of Willmar Manufacturing

X-77 is a registered trademark of Loveland Industries, Inc.