

# Minerva

## Fungicide

GROUP 3 FUNGICIDE

Application Type

AG

Agricultural

### ACTIVE INGREDIENT:

Tetraconazole {1-[2-(2,4-dichlorophenyl)-3-(1,1,2,2-tetrafluoroethoxy)

propyl]-1H-1,2,4-triazole} ..... 11.6%

**OTHER INGREDIENTS:** ..... 88.4%

**TOTAL** ..... 100.0%

Contains 1 lb. active ingredient (tetraconazole) per gallon.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### FIRST AID

<b>IF SWALLOWED</b>	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have affected person sip a glass of water if able to swallow.</li><li>• Do not induce vomiting unless told by a poison control center or doctor.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 858-7378 NPIC (human and animal health)

SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA Registration No. 60063-12

EPA Est. No. 70815-GA-001 (Lot No. begins with CB)

EPA Est. No. 60063-GA-001 (Lot No. begins with VL)

EPA Est. No. 86555-MO-001 (Lot No. begins with AF)

**MANUFACTURED FOR: SipcamAgro USA, Inc.**  
2525 Meridian Parkway, Suite 350  
Durham, NC 27713



SipcamAdvan

# **PRECAUTIONARY STATEMENTS**

## **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

### **CAUTION**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### **PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, polyvinyl chloride (PVC)  $\geq 14$  mils, and viton  $\geq 14$  mils. If you want more options, follow the instructions for category C on an EPA chemical-resistance category selection chart.

#### **Applicators and other handlers must wear:**

- Long sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### **USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### **ENVIRONMENTAL HAZARDS**

This product may be toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms adjacent to treatment areas. Exercise care when making applications of this product, and do not apply when atmospheric conditions favor drift or runoff. Do not contaminate water when disposing of equipment washwaters or rinsate.

In order to mitigate concern for reproductive effects to endangered bird and mammal species which may occur incidentally in sugarbeet growing areas, you are required to ascertain through the state Department of Agriculture, or Cooperative Extension Service, whether the treatment area may contain habitat of Federally listed bird and mammal species; if so, treatment must be avoided in these areas.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. For early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

### **GENERAL USE INFORMATION:**

Apply Minerva Fungicide in water carrier by spraying onto crop surfaces that are intended to be protected from disease. To obtain adequate coverage of typical agricultural crops, total spray volume usually will range from 20 to 150 gallons per acre (200 to 1400 liters per hectare) for dilute sprays, and 5 to 10 gallons per acre (50 to 100 liters per hectare) for concentrate ground sprays and aircraft applications. Both ground and aircraft methods of application are recommended.

**Minerva Fungicide** should be added to the spray tank while filling with water. Keep the agitator running when filling spray tank and during spray operations. When tank mixing this product with other pesticides observe the more restrictive label limitations and precautions. Do not exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing. Combination in the spray tank with other pesticides, fertilizers or surfactants is not recommended unless prior use has shown the combination to be physically compatible, effective and noninjurious under your conditions of use. When an adjuvant is to be used with this product, Sipcam Agro USA recommends the use of a Chemical Producers and Distributors Association (CPDA) certified adjuvant.

**Minerva Fungicide** contains a sterol demethylation inhibitor (sterol DMI) fungicide. For best results, apply this product in tank mix or in alternate applications with non-DMI fungicides as a resistance management strategy. Do not mix or alternate Minerva Fungicide with other DMI fungicides.

**AGRICULTURAL CROPS:**

Crop	Diseases Controlled	Minerva Fungicide Rate per Acre	Pre-harvest Interval	Application Directions
Sugarbeets	Cercospora leafspot, Powdery Mildew, Ramularia	13 fluid ounces	14 days	Apply Minerva Fungicide when conditions are favorable for Cercospora leafspot or powdery mildew. Do not make more than one (1) application of this product per growing season. Crops other than sugarbeet and small grains* must not be grown within 120 days following the last application of Minerva Fungicide. *Small grains can be planted 45 days following last application of this product.

**GENERAL CHEMIGATION INSTRUCTIONS:**

Apply this product only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation system. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

For specific information about calibration, contact State Extension Service specialists, equipment manufacturers, or other irrigation experts. Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

**Prevent the movement of Minerva Fungicide into the soil:**

- Minimize pesticide contact with the soil surface by chemigating above the crop canopy.
- Stop chemigation when pesticide mixture is observed running off crop surfaces or after 0.25 inches of water has been applied, whichever occurs first.
- Allow for sufficient time after chemigation for crop surfaces to dry prior to expected rainfall or to irrigation applied above the crop canopy.

**Sprinkler Chemigation:**

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

When mixing, fill nurse tank half full with water. Add Minerva Fungicide slowly to tank while hydraulic or mechanical agitation is operating and continue filling with water. Stickers, spreaders, etc., should be added last. If compatibility is in question, use the compatibility jar test before mixing a whole tank.

Because of the wide variety of possible combinations which can be encountered, observe all cautions and limitations on the label of all products used in mixtures.

Minerva Fungicide should be added through a traveling irrigation system continuously or at the last 30 minutes of solid set or hand moved irrigation systems. Agitation is recommended.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a dry, temperature-controlled, secure place.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** For 2.5 gal jugs: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**For 15 gal. and 30 gal. containers:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**FOR BULK AND MINI-BULK CONTAINERS** [4000 gal. tanker truck and 260 gal. totes]

**Container Disposal:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

When the container is empty, replace the cap and seal all openings that have been opened during use; and return to the point of purchase, or to a designated location named at the time of purchase of this product. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged or leaking, call Chem-Trec. If the container is damaged and leaking or material has been spilled, follow these procedures:

- Cover spill with absorbent material.
- Sweep into disposal container.
- Wash area with detergent and water and follow with clean water rinse.
- Do not allow to contaminate water supplies.
- Dispose of according to instructions.

If not returned to the point of purchase or to a designated location, clean empty container as instructed above and offer for recycling. Disposal of this container must be in compliance with state and local regulations.

**THIS CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.**

## WARRANTY AND LIMITATION OF DAMAGES

Conditions of sale: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.